

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|---|
| (51) International Patent Classification ⁶ : A61K 31/38, 31/40, 31/42, 31/165, 31/445, C07C 233/64, C07D 209/44, 211/32, 333/06 | | A1 | (11) International Publication Number: WO 99/00127 (43) International Publication Date: 7 January 1999 (07.01.99) |
| (21) International Application Number: PCT/US98/13424 (22) International Filing Date: 26 June 1998 (26.06.98) (30) Priority Data: 60/050,885 26 June 1997 (26.06.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BEIGHT, Douglas, Wade [US/US]; Apartment G, 9368 Benchmark Drive, Indianapo- lis, IN 46240 (US). CRAFT, Trelia, Joyce [US/US]; 10404 East 46th Street, Indianapolis, IN 46236 (US). FRAN- CISKOVICH, Jeffery, Bernard [US/US]; 5036 Quail Ridge Lane, Indianapolis, IN 46254 (US). GOODSON, Theodore, Junior [US/US]; 4045 Devon Drive, Indianapolis, IN 46226 (US). HALL, Steven, Edward [US/US]; 102 Nuttall Place, Chapel Hill, NC 27514 (US). HERRON, David, Kent [US/US]; 5945 Andover Road, Indianapolis, IN 46220 (US). KLIMKOWSKI, Valentine, Joseph [US/US]; 4504 Camelot Lane, Carmel, IN 46033 (US). MASTERS, John, Joseph [US/US]; 8338 Crystal Pointe Lane, Indianapolis, | | IN 46236 (US). MENDEL, David [US/US]; 11348 Woods Bay Lane, Indianapolis, IN 46236 (US). MILOT, Guy [CA/US]; 2 Farrington Street, Foxborough, MA 02035 (US). SAWYER, Jason, Scott [US/US]; 5718 North Winthrop Avenue, Indianapolis, IN 46220 (US). SHUMAN, Robert, Theodore [US/US]; 830 Ashton Park Drive, Greenwood, IN 46143 (US). SMITH, Gerald, Floyd [US/US]; 825 Queenswood Court, Indianapolis, IN 46217 (US). TEBBE, Anne, Louise [US/US]; 6202 North Sherman Drive, In- dianapolis, IN 46220 (US). TINSLEY, Jennifer, Marie [US/US]; 4542 State Road 39 North, Martinsville, IN 46151 (US). WEIR, Leonard, Crayton [US/US]; 6520 Englehardt Drive, Raleigh, NC 27613 (US). WIKEL, James, Howard [US/US]; 4068 Sunshine Way, Greenwood, IN 46142 (US). WILEY, Michael, Robert [US/US]; 7725 Langwood Drive, Indianapolis, IN 46268 (US). YEE, Ying, Kwong [US/US]; 5127 Briarstone Trace, Carmel, IN 46033 (US). (74) Agents: JACKSON, Thomas, E. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. | |
| (54) Title: ANTITHROMBOTIC AGENTS | | | |
| (57) Abstract | | | |
| This application relates to a compound of formula (I) (or a prodrug thereof or a pharmaceutically acceptable salt of the compound or prodrug thereof) as defined herein, pharmaceutical compositions thereof, and its use as an inhibitor of factor Xa, as well as a process for its preparation and intermediates therefor. | | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|---------------------|----|-----------------------|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LJ | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | Republic of Macedonia | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's | NZ | New Zealand | | |
| CM | Cameroon | | Republic of Korea | PL | Poland | | |
| CN | China | KR | Republic of Korea | PT | Portugal | | |
| CU | Cuba | KZ | Kazakhstan | RO | Romania | | |
| CZ | Czech Republic | LC | Saint Lucia | RU | Russian Federation | | |
| DE | Germany | LI | Liechtenstein | SD | Sudan | | |
| DK | Denmark | LK | Sri Lanka | SE | Sweden | | |
| EE | Estonia | LR | Liberia | SG | Singapore | | |

- 1 -

ANTITHROMBOTIC AGENTS

This application claims the benefit of U.S. Provisional
5 Application No. 60/050,885, filed June 26, 1997.

This invention relates to antithrombotic carboximid-
amide derivatives which demonstrate activity as inhibitors
of factor Xa and, accordingly, which are useful
anticoagulants in mammals. In particular it relates to
10 carboximidamide derivatives having high anticoagulant
activity, and antithrombotic activity. Thus, this invention
relates to new inhibitors of factor Xa, pharmaceutical
compositions containing the compounds as active ingredients,
and the use of the compounds as anticoagulants for
15 prophylaxis and treatment of thromboembolic disorders such
as venous thrombosis, pulmonary embolism, arterial
thrombosis, in particular myocardial ischemia, myocardial
infarction and cerebral thrombosis, general hypercoagulable
states and local hypercoagulable states, such as following
20 angioplasty and coronary bypass operations, and generalized
tissue injury as it relates to the inflammatory process. In
addition, the antithrombotic agents are useful as
anticoagulants in in vitro applications.

The process of blood coagulation, thrombosis, is
25 triggered by a complex proteolytic cascade leading to the
formation of thrombin. Thrombin proteolytically removes
activation peptides from the A α -chains and the B β -chains of
fibrinogen, which is soluble in blood plasma, initiating
insoluble fibrin formation. The formation of thrombin from
30 prothrombin is catalyzed by factor Xa.

- 2 -

Anticoagulation currently is achieved by the administration of heparins and coumarins. Parenteral pharmacological control of coagulation and thrombosis is based on inhibition of thrombin through the use of heparins.

5 Heparins act indirectly on thrombin by accelerating the inhibitory effect of endogenous antithrombin III (the main physiological inhibitor of thrombin). Because antithrombin III levels vary in plasma and because clot-bound thrombin seems resistant to this indirect mechanism, heparins can be

10 an ineffective treatment. Because coagulation assays are believed to be associated with efficacy and with safety, heparin levels must be monitored with coagulation assays (particularly the activated partial thromboplastin time (APTT) assay). Coumarins impede the generation of thrombin

15 by blocking the posttranslational gamma-carboxylation in the synthesis of prothrombin and other proteins of this type. Because of their mechanism of action, the effect of coumarins can only develop slowly, 6-24 hours after administration. Further, they are not selective

20 anticoagulants. Coumarins also require monitoring with coagulation assays (particularly the prothrombin time (PT) assay).

Recently, interest has grown in small synthetic molecules which demonstrate potent direct inhibition of

25 thrombin and factor Xa. See, Jeremy J. Edmunds and Stephen T. Rapundalo (Annette M. Doherty, Section Editor), Annual Reports in Medicinal Chemistry, (1996), 31, 51-60.

Although the heparins and coumarins are effective anticoagulants, no commercial drug has yet emerged from the


30 small synthetic molecules; and despite the continuing promise for this class of compounds, there still exists a need for anticoagulants which act selectively on factor Xa or thrombin, and which, independent of antithrombin III, exert inhibitory action shortly after administration,

- 3 -

preferably by an oral route, and do not interfere with lysis of blood clots, as required to maintain hemostasis.

The present invention is directed to the discovery that the compounds of the present invention, as defined below,
 5 are potent inhibitors of factor Xa which may have high bioavailability following oral administration.

According to the invention there is provided a method of inhibiting factor Xa comprising using an effective amount of a factor Xa inhibiting compound of formula I
 10



wherein

A³, A⁴, A⁵ and A⁶, together with the two carbons to which they are attached, complete a substituted benzene in
 15 which A³ is CR³, A⁴ is CR⁴, A⁵ is CR⁵, and A⁶ is CR⁶;
 wherein

R³ is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy (which may bear an ω-carboxy substituent), benzoyloxy (which may bear one or more halo, hydroxy methoxy or methyl
 20 substitutents), methyl or methoxy;

one of R⁴ and R⁵ is hydrogen, methyl, halo, trifluoromethyl, nitro, amino(imino)methyl, amino(hydroxyimino)-methyl, R^fO-, R^fO₂C-, R^fO₂C-CH₂-, R^fO₂C-CH₂-O-, 3-methoxycarbonyl-1-oxopropyl, R^gNH- or bis(methylsulfonyl)-
 25 amino;

the other of R⁴ and R⁵ is hydrogen, halo or methyl; and
 R⁶ is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy (which may bear an ω-carboxy substituent), benzoyloxy (which may bear one or more halo, hydroxy methoxy or methyl
 30 substitutents), methyl or methoxy;

in which R^f is hydrogen, (1-4C)alkyl or benzyl; R^g is hydrogen, acetyl, trifluoroacetyl, phenylalanyl,

- 4 -

2-(t-butoxycarbonylamino)-4-methylsulfinyl-1-oxobutyl or R^hSO_2- ; and R^h is (1-4C)alkyl, trifluoromethyl, phenyl, 3,5-dimethylisoxazol-4-yl or dimethylamino; or

two adjacent residues selected from R^3 , R^4 , R^5 and R^6 together form a benz ring; and the other two are each hydrogen; or

A^3 , A^4 , A^5 and A^6 , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which

- 10 (a) one of A^3 , A^4 , A^5 and A^6 is N, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively;
- (b) two adjacent residues of A^3 , A^4 , A^5 and A^6 together form S, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively;
- 15 (c) two non-adjacent residues of A^3 , A^4 , A^5 and A^6 are each N, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively; or
- (d) A^3 and A^4 together form a fused benz ring, and A^5 and A^6 together form -NH-;


20 wherein

each of R^3 , R^4 , R^5 and R^6 is hydrogen, or one or two of R^3 , R^4 , R^5 and R^6 is independently chloro, bromo or methyl and the others are hydrogen;

L^1 is -NH-CO-, -CO-NH- or -O-CH₂- such that - L^1 - Q^1 is -NH-CO- Q^1 , -CO-NH- Q^1 or -O-CH₂- Q^1 (provided that when - L^1 - Q^1 is -CO-NH- Q^1 , then R^2 is not -CO-NH- Q^1 or -CO-NH- Q^{2B});

Q^1 is Q^{1A} , Q^{1B} or Q^{1C} wherein

Q^{1A} is




30

- 5 -


in which R^a is hydrogen, R^b is hydrogen and R^c is hydrogen or hydroxy; or R^a is hydroxy, R^b is hydrogen and R^c is hydrogen; or R^a and R^b together form a methylene or oxo group and R^c is hydrogen;

5 Q^{1B} is




in which R^d is hydrogen or hydroxy and R^e is hydrogen or fluoro; and

10 Q^{1C} is



R^2 is a residue defined for $-NH-CO-Q^1$ or $-CO-NH-Q^1$; or R^2 is $-L^{2A}-Q^{2A}$, $-L^{2B}-Q^{2B}$, $-L^{2C}-Q^{2C}$ or $-L^{2D}-Q^{2D}$ wherein L^{2A} is a direct bond; and

15 Q^{2A} is




in which D is carbonyl or $-CHR^k-$ in which R^k is hydrogen, hydroxy, (1-6C)alkoxy or $-CH_2-R^j$ in which R^j is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of R^m and R^n is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or R^m and R^n together form a

25 benz ring;

- 6 -

L^{2B} is -NH-CO-, -CO-NH-, -O-CO-, -CH₂-O- or -O-CH₂-
 such that -L^{2B}-Q^{2B} is -NH-CO-Q^{2B}, -CO-NH-Q^{2B}, -O-CO-Q^{2B},
 -CH₂-O-Q^{2B} or -O-CH₂-Q^{2B}; and
 Q^{2B} is



5

- in which R^O is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and R^P is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl,
- 10 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or -J-R^Q in which J is a single bond, methylene, carbonyl, oxo, -S(O)_q- (wherein q is 0, 1 or 2), or -NR^r- (wherein R^r is hydrogen or methyl); and R^Q is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;
- 15 L^{2C} is -NR^V-CO-X-, -NR^V-CS-Y-, -CH₂-CO-NR^W-CH₂-, -O-CO-, -O-CH₂-, -S-CH₂- or -CH₂-NR^X-CH₂- such that -L^{2C}-Q^{2C} is -NR^V-CO-X-Q^{2C}, -NR^V-CS-Y-Q^{2C}, -CH₂-CO-NR^W-CH₂-Q^{2C}, -O-CO-Q^{2C}, -O-CH₂-Q^{2C}, -S-CH₂-Q^{2C} or -CH₂-NR^X-CH₂-Q^{2C} in which X is -(CH₂)_x- (wherein x is 0, 1 or 2), -NR^W-CH₂-,
- 20 -O-CH₂- or -S-CH₂-; Y is -NR^W-CH₂- or -O-CH₂-; each of R^V and R^W is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the α-position; and R^X is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and
- Q^{2C} is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl
- 25 may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;
- L^{2D} is -NH-CO- such that -L^{2D}-Q^{2D} is -NH-CO-Q^{2D}; and
- Q^{2D} is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a
- 30 chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a

- 7 -

substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and $-\text{CH}_2-\text{R}^Z$ in which R^Z is isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents

5 independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent;

or a prodrug of the compound of formula I;

10 or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

In addition, there is provided the use of a factor Xa inhibiting compound of formula I (or prodrug or salt) as described herein as an active ingredient in the manufacture
15 of a medicament for use in producing an anticoagulant or antithrombotic effect.

The present invention also provides a method of inhibiting coagulation in a mammal comprising administering to a mammal in need of treatment, a coagulation inhibiting
20 dose of a factor Xa inhibiting compound of formula I having any of the definitions herein.

The present invention further provides a method of inhibiting factor Xa comprising administering to a mammal in need of treatment, a factor Xa inhibiting dose of a
25 factor Xa inhibiting compound of formula I having any of the definitions herein.

Further, the present invention provides a method of treating a thromboembolic disorder comprising administering to a mammal in need of treatment, an effective dose of a
30 factor Xa inhibiting compound of formula I having any of the definitions herein.

In addition, there is provided the use of a factor Xa inhibiting compound of formula I having any of the

- 8 -

definitions herein for the manufacture of a medicament for treatment of a thromboembolic disorder.

As an additional feature of the invention there is provided a pharmaceutical formulation comprising in
5 association with a pharmaceutically acceptable carrier, diluent or excipient, a prodrug of a factor Xa inhibiting compound of formula I (or of a pharmaceutically acceptable salt thereof) as provided in any of the descriptions herein.

In general, the factor Xa inhibiting compounds of
10 formula I are believed to be novel and, thus, to constitute an additional aspect of the invention. Thus, according to the invention there is provided a novel compound of formula I (or a pharmaceutically acceptable salt thereof) according to any of the definitions herein of a compound of
15 formula I, provided that the compound is not one which is not novel.

A pharmaceutically acceptable salt of an antithrombotic agent of the instant invention includes one which is an acid-addition salt made from a basic compound of formula I
20 and an acid which provides a pharmaceutically acceptable anion, as well as a salt which is made from an acidic compound of formula I and a base which provides a pharmaceutically acceptable cation. Thus, a salt of a novel compound of formula I as provided herein made with an acid
25 or base which affords a pharmaceutically acceptable counterion provides a particular aspect of the invention. Examples of such acids and bases are provided hereinbelow.

As an additional aspect of the invention there is provided a pharmaceutical formulation comprising in
30 association with a pharmaceutically acceptable carrier, diluent or excipient, a novel compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in any of the descriptions herein.

- 9 -

In this specification, the following definitions are used, unless otherwise described: Halo is fluoro, chloro, bromo or iodo. Alkyl, alkoxy, etc. denote both straight and branched groups; but reference to an individual radical such as "propyl" embraces only the straight chain ("normal") radical, a branched chain isomer such as "isopropyl" being specifically denoted. When two adjacent residues form a (fused) benz ring, they form a cis,cis-buta-1,3-dien-1,4-diyl divalent radical.

It will be appreciated that certain compounds of formula I (or salts or prodrugs, etc.) may exist in, and be isolated in, isomeric forms, including tautomeric forms, cis- or trans-isomers, as well as optically active, racemic, or diastereomeric forms. It is to be understood that the present invention encompasses a compound of formula I in any of the tautomeric forms or as an a mixture thereof; or as a mixture of diastereomers, as well as in the form of an individual diastereomer, and that the present invention encompasses a compound of formula I as a mixture of enantiomers, as well as in the form of an individual enantiomer, any of which mixtures or form possesses inhibitory properties against factor Xa, it being well known in the art how to prepare or isolate particular forms and how to determine inhibitory properties against factor Xa by standard tests including those described below.


In addition, a compound of formula I (or salt or prodrug, etc.) may exhibit polymorphism or may form a solvate with water or an organic solvent. The present invention also encompasses any such polymorphic form, any solvate or any mixture thereof.

Particular values are listed below for radicals, substituents, and ranges, for illustration only, and they do not exclude other defined values or other values within defined ranges for the radicals and substituents.

- 10 -

For an alkyl group or the alkyl portion of an alkyl containing group such as, for example alkoxy, a particular value for (1-2C)alkyl is methyl or ethyl, and more particularly is methyl; for (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl, and more particularly is methyl, isopropyl, butyl or t-butyl; for (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or hexyl, and more particularly is methyl, butyl, or hexyl. A particular value for halo is bromo or chloro, and more particularly is chloro.

A particular compound of formula I is one of formula Ia



wherein A⁴, L¹, Q¹ and R² have any of the values defined herein.

A particular value for R² is, for example, (4-isopropylbenzoyl)amino, (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)amino, or [1-(4-pyridyl)piperidin-4-yl]-methoxycarbonylamino.

A particular value for R⁴ or R⁵ is, for example, hydrogen, hydroxy, acetylamino, carboxymethoxy, carboxymethyl, methoxycarbonyl or carboxy.

One particular compound of formula I as described herein is one in which L¹-Q¹ is -NH-CO-Q¹.

Another particular compound of formula I as described herein is one in which L¹-Q¹ is -CO-NH-Q¹.

A prodrug of a compound of formula I may be one formed in a conventional manner with a functional group of the compound, such as with an amino, hydroxy or carboxy group.

One particular prodrug is a carbamate in which an amino (or imino) group of Q¹ is substituted by a [(1-4C)alkoxy]-


- 11 -

carbonyl or acetoxymethoxycarbonyl group, for example as described in Examples 26-30.

A compound of formula I may be prepared by processes which include processes known in the chemical art for the production of any known compounds of formula I or of structurally analogous compounds or by a novel process described herein. A process for the preparation of a novel compound of formula I (or a pharmaceutically acceptable salt thereof), novel processes for the preparation of a compound of formula I and novel intermediates for the manufacture of a compound of formula I as defined above provide further features of the invention and are illustrated by the following procedures in which the meanings of the generic radicals are as defined above, unless otherwise specified. It will be recognized that it may be preferred or necessary to prepare a compound of formula I in which a functional group is protected using a conventional protecting group, then to remove the protecting group to provide the compound of formula I.

Thus, there is provided a process for preparing a novel compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in any of the above descriptions which is selected from any of those described in the examples, including the following.

(A) For a compound of formula I in which the linkage of R² to the ring terminates in -NH-CO-, -NR^V-CO- or -NR^V-CS-, acylating an amine of formula II,



II


30

or a corresponding amine in which the nitrogen bears the group R^V, using a corresponding acid which terminates with

- 12 -

the group HO-CO- or HO-CS-, or an activated derivative thereof. Typical activated derivatives include the acid halides, activated esters, including 4-nitrophenyl esters and those derived from coupling reagents, as well as (when
 5 the product is a urea or thiourea) isocyanates and isothiocyanates.

(B) For a compound of formula I in which $-L^1-Q^1$ is $-NH-CO-Q^1$, acylating an amine of formula III




III

10

using an acid of formula $HO-CO-Q^1$, or an activated derivative thereof.

(C) For a compound of formula I in which $-L^1-Q^1$ is
 15 $-CO-NH-Q^1$ and R^2 is of the form $-NH-CO-Q^2$, acylating an amine of formula H_2N-Q^1 using a [1,3]oxazine of formula IV,




IV

20 wherein Q^2 represents, for example, Q^{2B} , Q^{2C} or Q^{2D} .

(D) For a compound of formula I in which R^2 is $-L^{2A}-Q^{2A}$ and D is carbonyl, diacylating a compound of formula II using an anhydride of formula V.

- 13 -



(E) For a compound of formula I in which Q¹ is Q^{1A} and R^a is hydroxy or in which Q¹ is Q^{1B} and R^d is hydroxy,
 5 adding hydroxylamine to a corresponding nitrile to afford the compound with an amino(hydroxyimino)methyl group. Conveniently, the addition is carried out as described in Example 1, Part F.

(F) For a compound of formula I in which Q¹ is Q^{1A} and R^a, R^b and R^c are each hydrogen, or in which Q¹ is Q^{1B} and R^d is hydrogen, hydrogenolyzing the N-O bond of a
 10 corresponding compound of formula I in which R^a is hydroxy or in which R^d is hydroxy. The hydrogenolysis conveniently is carried out as described in Example 1, Part G.

(G) For a compound of formula I in which Q¹ is Q^{1A} and R^a, R^b and R^c are each hydrogen, or in which Q¹ is Q^{1B} and R^d is hydrogen, substituting the methylthio group of a
 15 corresponding compound bearing a -C(SCH₃)=NH group with an amino group by treatment with ammonia, or a solvate or salt thereof. Conveniently, the substitution is carried out
 20 using ammonium acetate as described in Example 15.

(H) For a compound of formula I in which Q¹ is Q^{1A} and R^a and R^b together form a methylene or oxo group, cyclizing a corresponding 4-aminomethyl-3-cyanophenyl or 4-aminooxo-
 25 3-cyanophenyl compound. Conveniently, the amine is generated and cyclized in situ, for example as described in Example 8, Part M or in Example 39.

(I) For a compound of formula I in which Q¹ is Q^{1A} and R^c is hydroxy, hydrogenolyzing the N-O bond of a
 30 corresponding 3-amino-1,2-benzisoxazol-7-yl compound.

- 14 -

Conveniently, the hydrogenolysis is carried out as described in Example 31.

Whereafter, for any of the above procedures, when a functional group is protected using a protecting group,
5 removing the protecting group.

Whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of a compound of formula I is required, it is obtained by reacting the basic form of a basic compound of formula I with an acid affording a
10 physiologically acceptable counterion or the acidic form of an acidic compound of formula I with a base affording a physiologically acceptable counterion or by any other conventional procedure.

A novel intermediate or starting material compound such
15 as, for example, a novel compound of formula II, III or IV, etc., provides a further aspect of the invention.

As mentioned above, a compound corresponding to a compound of formula I but in which a functional group is protected may serve as an intermediate for a compound of
20 formula I. Accordingly, such a protected intermediate for a novel compound of formula I provides a further aspect of the invention. Thus, as one particular aspect of the invention, there is provided a compound corresponding to a novel compound of formula I as defined above in which R⁴ which is
25 hydroxy, but in which the corresponding substituent is -OPP in place of hydroxy, wherein PP is a phenol protecting group other than (1-4C)alkyl or benzyl. Phenol protecting groups are well known in the art, for example as described in T.W. Greene and P.G.M. Wuts, "Protecting Groups in Organic
30 Synthesis" (1991). Further, PP may denote a functionalized resin, for example as disclosed in H.V. Meyers, et al., Molecular Diversity, (1995), 1, 13-20.

As mentioned above, the invention includes a pharmaceutically acceptable salt of the factor Xa inhibiting

- 15 -

compound defined by the above formula I. A basic compound of this invention possesses one or more functional groups sufficiently basic to react with any of a number of inorganic and organic acids affording a physiologically acceptable counterion to form a pharmaceutically acceptable salt. Acids commonly employed to form pharmaceutically acceptable acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromobenzenesulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts thus are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like. Preferred pharmaceutically acceptable acid addition salts include those formed with mineral acids such as hydrochloric acid, hydrobromic acid and sulfuric acid.

For a compound of formula I which bears an acidic moiety, such as a carboxy group, a pharmaceutically acceptable salt may be made with a base which affords a pharmaceutically acceptable cation, which includes alkali

- 16 -

metal salts (especially sodium and potassium), alkaline earth metal salts (especially calcium and magnesium), aluminum salts and ammonium salts, as well as salts made from physiologically acceptable organic bases such as triethylamine, morpholine, piperidine and triethanolamine.

If not commercially available, a necessary starting material for the preparation of a compound of formula I may be prepared by a procedure which is selected from standard techniques of organic chemistry, including aromatic and heteroaromatic substitution and transformation, from techniques which are analogous to the syntheses of known, structurally similar compounds, and techniques which are analogous to the above described procedures or procedures described in the Examples. It will be clear to one skilled in the art that a variety of sequences is available for the preparation of the starting materials. Starting materials which are novel provide another aspect of the invention.

Selective methods of substitution, protection and deprotection are well known in the art for preparation of a compound such as one of formula II, III, IV or VI discussed above.

Generally, a basic compound of the invention is isolated best in the form of an acid addition salt. A salt of a compound of formula I formed with an acid such as one of those mentioned above is useful as a pharmaceutically acceptable salt for administration of the antithrombotic agent and for preparation of a formulation of the agent. Other acid addition salts may be prepared and used in the isolation and purification of the compounds.

As noted above, the optically active isomers and diastereomers of the compounds of formula I are also considered part of this invention. Such optically active isomers may be prepared from their respective optically active precursors by the procedures described above, or by

- 17 -

resolving the racemic mixtures. This resolution can be carried out by derivatization with a chiral reagent followed by chromatography or by repeated crystallization. Removal of the chiral auxiliary by standard methods affords
5 substantially optically pure isomers of the compounds of the present invention or their precursors. Further details regarding resolutions can be obtained in Jacques, et al., Enantiomers, Racemates, and Resolutions, John Wiley & Sons, 1981.

10 The compounds of the invention are believed to selectively inhibit factor Xa over other proteinases and nonenzyme proteins involved in blood coagulation without appreciable interference with the body's natural clot lysing
15 fibrinolysis). Further, such selectivity is believed to permit use with thrombolytic agents without substantial interference with thrombolysis and fibrinolysis.

The invention in one of its aspects provides a method of inhibiting factor Xa in mammals comprising administering
20 to a mammal in need of treatment an effective (factor Xa inhibiting) dose of a compound of formula I.

In another of its aspects, the invention provides a method of treating a thromboembolic disorder comprising administering to a mammal in need of treatment an effective
25 (thromboembolic disorder therapeutic and/or prophylactic amount) dose of a compound of formula I.

The invention in another of its aspects provides a method of inhibiting coagulation in a mammal comprising administering to a mammal in need of treatment an effective
30 (coagulation inhibiting) dose of a compound of formula I.

The factor Xa inhibition, coagulation inhibition and thromboembolic disorder treatment contemplated by the present method includes both medical therapeutic and/or prophylactic treatment as appropriate.

- 18 -

In a further embodiment the invention relates to treatment, in a human or animal, of a condition where inhibition of factor Xa is required. The compounds of the invention are expected to be useful in mammals, including
5 man, in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disorders in which the compounds have a potential utility are in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disorders in which the compounds have a
10 potential utility, in treatment and/or prophylaxis, include venous thrombosis and pulmonary embolism, arterial thrombosis, such as in myocardial ischemia, myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial thrombosis. Further, the compounds have
15 expected utility in the treatment or prophylaxis of atherosclerotic disorders (diseases) such as coronary arterial disease, cerebral arterial disease and peripheral arterial disease. Further, the compounds are expected to be useful together with thrombolytics in myocardial infarction.
20 Further, the compounds have expected utility in prophylaxis for reocclusion after thrombolysis, percutaneous transluminal angioplasty (PTCA) and coronary bypass operations. Further, the compounds have expected utility in prevention of rethrombosis after microsurgery. Further, the
25 compounds are expected to be useful in anticoagulant treatment in connection with artificial organs and cardiac valves. Further, the compounds have expected utility in anticoagulant treatment in hemodialysis and disseminated intravascular coagulation. A further expected utility is in
30 rinsing of catheters and mechanical devices used in patients in vivo, and as an anticoagulant for preservation of blood, plasma and other blood products in vitro. Still further, the compounds have expected utility in other diseases where blood coagulation could be a fundamental contributing

- 19 -

process or a source of secondary pathology, such as cancer, including metastasis, inflammatory diseases, including arthritis, and diabetes. The anti-coagulant compound is administered orally or parenterally, e.g. by intravenous
5 infusion (iv), intramuscular injection (im) or subcutaneously (sc).

The specific dose of a compound administered according to this invention to obtain therapeutic and/or prophylactic effects will, of course, be determined by the particular
10 circumstances surrounding the case, including, for example, the compound administered, the rate of administration, the route of administration, and the condition being treated.

A typical daily dose for each of the above utilities is between about 0.01 mg/kg and about 1000 mg/kg. The dose
15 regimen may vary e.g. for prophylactic use a single daily dose may be administered or multiple doses such as 3 or 5 times daily may be appropriate. In critical care situations a compound of the invention is administered by iv infusion at a rate between about 0.01 mg/kg/h and about 20 mg/kg/h
20 and preferably between about 0.1 mg/kg/h and about 5 mg/kg/h.

The method of this invention also is practiced in conjunction with a clot lysing agent e.g. tissue plasminogen activator (t-PA), modified t-PA, streptokinase or urokinase.
25 In cases when clot formation has occurred and an artery or vein is blocked, either partially or totally, a clot lysing agent is usually employed. A compound of the invention can be administered prior to or along with the lysing agent or subsequent to its use, and preferably further is
30 administered along with aspirin to prevent the reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with a platelet glycoprotein receptor (IIb/IIIa) antagonist, that inhibits platelet aggregation. A compound

- 20 -

of the invention can be administered prior to or along with the IIb/IIIa antagonist or subsequent to its use to prevent the occurrence or reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with aspirin. A compound of the invention can be administered prior to or along with aspirin or subsequent to its use to prevent the occurrence or reoccurrence of clot formation. As stated above, preferably a compound of the present invention is administered in conjunction with a clot lysing agent and aspirin.

This invention also provides a pharmaceutical composition for use in the above described therapeutic method. A pharmaceutical composition of the invention comprises an effective factor Xa inhibiting amount of a compound of formula I in association with a pharmaceutically acceptable carrier, excipient or diluent.

The active ingredient in such formulations comprises from 0.1 percent to 99.9 percent by weight of the formulation. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

For oral administration the antithrombotic compound is formulated in gelatin capsules or tablets which may contain excipients such as binders, lubricants, disintegration agents and the like. For parenteral administration the antithrombotic is formulated in a pharmaceutically acceptable diluent e.g. physiological saline (0.9 percent), 5 percent dextrose, Ringer's solution and the like.

The compound of the present invention can be formulated in unit dosage formulations comprising a dose between about 0.1 mg and about 1000 mg. Preferably the compound is in the form of a pharmaceutically acceptable salt such as for example the sulfate salt, acetate salt or a phosphate salt.

- 21 -

An example of a unit dosage formulation comprises 5 mg of a compound of the present invention as a pharmaceutically acceptable salt in a 10 mL sterile glass ampoule. Another example of a unit dosage formulation comprises about 10 mg
5 of a compound of the present invention as a pharmaceutically acceptable salt in 20 mL of isotonic saline contained in a sterile ampoule.

The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous,
10 intravenous, intramuscular, and intranasal. The compounds of the present invention are preferably formulated prior to administration.

The present pharmaceutical compositions are prepared by known procedures using well known and readily available
15 ingredients. The compositions of this invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art. In making the compositions of the present invention, the active
20 ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a
25 vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, (as a solid or in a liquid medium), soft and hard gelatin capsules,
30 suppositories, sterile injectable solutions, sterile packaged powders, and the like.

The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way. "Active ingredient," of course, means

- 22 -

a compound according to formula I or a pharmaceutically acceptable salt or solvate thereof.

- Formulation 1: Hard gelatin capsules are prepared
5 using the following ingredients:

| | Quantity (mg/capsule) |
|--------------------|--------------------------|
| Active ingredient | 250 |
| Starch, dried | 200 |
| Magnesium stearate | <u>10</u> |
| Total | 460 mg |

- Formulation 2: A tablet is prepared using the
ingredients below:

| | Quantity (mg/tablet) |
|-----------------------------|-------------------------|
| Active ingredient | 250 |
| Cellulose, microcrystalline | 400 |
| Silicon dioxide, fumed | 10 |
| Stearic acid | <u>5</u> |
| Total | 665 mg |

- 10 The components are blended and compressed to form tablets
each weighing 665 mg.

- Formulation 3: An aerosol solution is prepared
containing the following components:

| | Weight |
|---------------------------------------|--------------|
| Active ingredient | 0.25 |
| Ethanol | 29.75 |
| Propellant 22 (Chlorodifluoromethane) | <u>70.00</u> |
| Total | 100.00 |

- 23 -

The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

Formulation 4: Tablets, each containing 60 mg of active ingredient, are made as follows:

| | |
|---|-------------|
| Active ingredient | 60 mg |
| Starch | 45 mg |
| Microcrystalline cellulose | 35 mg |
| Polyvinylpyrrolidone (as 10% solution in water) | 4 mg |
| Sodium carboxymethyl starch | 4.5 mg |
| Magnesium stearate | 0.5 mg |
| Talc | <u>1 mg</u> |
| Total | 150 mg |

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50 °C and passed through a No. 18 mesh U.S. Sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

- 24 -

Formulation 5: Capsules, each containing 80 mg of active ingredient, are made as follows:

| | |
|----------------------------|-------------|
| Active ingredient | 80 mg |
| Starch | 59 mg |
| Microcrystalline cellulose | 59 mg |
| Magnesium stearate | <u>2 mg</u> |
| Total | 200 mg |

- 5 The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

- 10 Formulation 6: Suppositories, each containing 225 mg of active ingredient, are made as follows:

| | |
|---------------------------------|-----------------|
| Active ingredient | 225 mg |
| Saturated fatty acid glycerides | <u>2,000 mg</u> |
| Total | 2,225 mg |

- 15 The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

- 20 Formulation 7: Suspensions, each containing 50 mg of active ingredient per 5 mL dose, are made as follows:

| | |
|--------------------------------|---------|
| Active ingredient | 50 mg |
| Sodium carboxymethyl cellulose | 50 mg |
| Syrup | 1.25 mL |
| Benzoic acid solution | 0.10 mL |

- 25 -

| | |
|-------------------------|------|
| Flavor | q.v. |
| Color | q.v. |
| Purified water to total | 5 mL |

The active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8: An intravenous formulation may be prepared as follows:

| | |
|-------------------|----------|
| Active ingredient | 100 mg |
| Isotonic saline | 1,000 mL |

The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 mL per minute.

The ability of a compound of the present invention to be an effective and orally active factor Xa inhibitor may be evaluated in one or more of the following assays or in other standard assays known to those in the art.

The inhibition by a compound of the inhibition of a serine protease of the human blood coagulation system or of the fibrinolytic system, as well as of trypsin, is determined in vitro for the particular enzyme by measuring its inhibitor binding affinity in an assay in which the enzyme hydrolyzes a particular chromogenic substrate, for example as described in Smith, G.F.; Gifford-Moore, D.; Craft, T.J.; Chirgadze, N.; Ruterbories, K.J.; Lindstrom, T.D.; Satterwhite, J.H. Efegatran: A New Cardiovascular Anticoagulant. *New Anticoagulants for the Cardiovascular*

- 26 -

Patient; Pifarre, R., Ed.; Hanley & Belfus, Inc.:
Philadelphia, 1997; pp. 265-300. The inhibitor binding
affinity is measured as apparent association constant K_{ass}
which is the hypothetical equilibrium constant for the
5 reaction between enzyme and the test inhibitor compound (I).



$$K_{ass} = \frac{[\text{Enzyme-I}]}{[(\text{Enzyme}) \times (\text{I})]}$$

Conveniently, enzyme inhibition kinetics are performed
10 in 96-well polystyrene plates and reaction rates are
determined from the rate of hydrolysis of appropriate
p-nitroanilide substrates at 405 nm using a Thermomax plate
reader from Molecular Devices (San Francisco, CA). The same
protocol is followed for all enzymes studied: 50 μL buffer
15 (0.03 M Tris, 0.15 M NaCl pH 7) in each well, followed by
25 μL of inhibitor solution (in 100% methanol, or in 50% v:v
aqueous methanol) and 25 μL enzyme solution; within two
minutes, 150 μL aqueous solution of chromogenic substrate
(0.25 mg/mL) is added to start the enzymatic reaction. The
20 rates of chromogenic substrate hydrolysis reactions provide
a linear relationship with the enzymes studied such that
free enzyme can be quantitated in reaction mixtures. Data
is analyzed directly as rates by the Softmax program to
produce [free enzyme] calculations for tight-binding K_{ass}
25 determinations. For apparent K_{ass} determinations, 1.34 nM
human factor Xa is used to hydrolyze 0.18 mM BzIle-Glu-Gly-
Arg-pNA; 5.9 nM human thrombin or 1.4 nM bovine trypsin is
used to hydrolyze 0.2 mM BzPhe-Val-Arg-pNA; 3.4 nM human
plasmin is used with 0.5 mM HD-Val-Leu-Lys-pNA; 1.2 nM human
30 nt-PA is used with 0.81 mM HD-Ile-Pro-Arg-pNA; and 0.37 nM
urokinase is used with 0.30 mM pyro-gfsGlu-Gly-Arg-pNA.

- 27 -

Kass is calculated for a range of concentrations of test compounds and the mean value reported in units of liter per mole. In general, a factor Xa inhibiting compound of formula I of the instant invention exhibits a Kass of 0.1 to 5 0.5 x 10⁶ L/mole or much greater.

The factor Xa inhibitor preferably should spare fibrinolysis induced by urokinase, tissue plasminogen activator (t-PA) and streptokinase. This would be important to the therapeutic use of such an agent as an adjunct to 10 streptokinase, tp-PA or urokinase thrombolytic therapy and to the use of such an agent as an endogenous fibrinolysis-sparing (with respect to t-PA and urokinase) antithrombotic agent. In addition to the lack of interference with the amidase activity of the fibrinolytic proteases, such 15 fibrinolytic system sparing can be studied by the use of human plasma clots and their lysis by the respective fibrinolytic plasminogen activators.

Materials

20 Dog plasma is obtained from conscious mixed-breed hounds (either sex Butler Farms, Clyde, New York, U.S.A.) by venipuncture into 3.8 percent citrate. Fibrinogen is prepared from fresh dog plasma and human fibrinogen is prepared from in-date ACD human blood at the fraction I-2 25 according to previous procedures and specification. Smith, Biochem. J., 185, 1-11 (1980; and Smith, et al., Biochemistry, 11, 2958-2967, (1972). Human fibrinogen (98 percent pure/plasmin free) is from American Diagnostica, Greenwich, Connecticut. Radiolabeling of fibrinogen I-2 30 preparations is performed as previously reported. Smith, et al., Biochemistry, 11, 2958-2967, (1972). Urokinase is purchased from Leo Pharmaceuticals, Denmark, as 2200 Ploug units/vial. Streptokinase is purchased from Hoechst-Roussel Pharmaceuticals, Somerville, New Jersey.

- 28 -

Methods - Effects on Lysis of Human Plasma Clots by t-PA

Human plasma clots are formed in micro test tubes by adding 50 μ L thrombin (73 NIH unit/mL) to 100 μ L human plasma which contains 0.0229 μ Ci 125-iodine labeled fibrinogen. Clot lysis is studied by overlaying the clots with 50 μ L of urokinase or streptokinase (50, 100, or 1000 unit/mL) and incubating for 20 hours at room temperature. After incubation the tubes are centrifuged in a Beckman Microfuge. 25 μ L of supernate is added into 1.0 mL volume of 0.03 M tris/0.15 M NaCl buffer for gamma counting. Counting controls 100 percent lysis are obtained by omitting thrombin (and substituting buffer). The factor Xa inhibitors are evaluated for possible interference with fibrinolysis by including the compounds in the overlay solutions at 1, 5, and 10 μ g/mL concentrations. Rough approximations of IC₅₀ values are estimated by linear extrapolations from data points to a value which would represent 50 percent of lysis for that particular concentration of fibrinolytic agent.

Anticoagulant ActivityMaterials

Dog plasma and rat plasma are obtained from conscious mixed-breed hounds (either sex, Butler Farms, Clyde, New York, U.S.A.) or from anesthetized male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc., Indianapolis, Indiana, U.S.A.) by venipuncture into 3.8 percent citrate. Fibrinogen is prepared from in-date ACD human blood as the fraction I-2 according to previous procedures and specifications. Smith, Biochem. J., 185, 1-11 (1980); and Smith, et al., Biochemistry, 11, 2958-2967 (1972). Human fibrinogen is also purchased as 98 percent pure/plasmin free from American Diagnostica, Greenwich, Connecticut. Coagulation reagents Actin, Thromboplastin, Innovin and Human plasma are from Baxter Healthcare Corp., Dade Division, Miami, Florida.

- 29 -

Bovine thrombin from Parke-Davis (Detroit, Michigan) is used for coagulation assays in plasma.

Methods

5 Anticoagulation Determinations

Coagulation assay procedures are as previously described. Smith, et al., Thrombosis Research, 50, 163-174 (1988). A CoAScreener coagulation instrument (American LABor, Inc.) is used for all coagulation assay measurements. The
10 prothrombin time (PT) is measured by adding 0.05 mL saline and 0.05 mL Thromboplastin-C reagent or recombinant human tissue factor reagent (Innovin) to 0.05 mL test plasma. The activated partial thromboplastin time (APTT) is measured by incubation of 0.05 mL test plasma with 0.05 mL Actin reagent
15 for 120 seconds followed by 0.05 mL CaCl₂ (0.02 M). The thrombin time (TT) is measured by adding 0.05 mL saline and 0.05 mL thrombin (10 NIH units/mL) to 0.05 mL test plasma. The compounds of formula I are added to human or animal plasma over a wide range of concentrations to determine
20 prolongation effects on the APTT, PT, and TT assays. Linear extrapolations are performed to estimate the concentrations required to double the clotting time for each assay.

Animals

25 Male Sprague Dawley rats (350-425 gm, Harlan Sprague Dawley Inc., Indianapolis, IN) are anesthetized with xylazine (20-mg/kg, s.c.) and ketamine (120 mg/kg, s.c.) and maintained on a heated water blanket (37 °C). The jugular vein(s) is cannulated to allow for infusions.

30

Arterio-Venous shunt model

The left jugular vein and right carotid artery are cannulated with 20 cm lengths of polyethylene PE 60 tubing. A 6 cm center section of larger tubing (PE 190) with a
35 cotton thread (5 cm) in the lumen, is friction fitted between the longer sections to complete the arterio-venous

- 30 -

shunt circuit. Blood is circulated through the shunt for 15 min before the thread is carefully removed and weighed. The weight of a wet thread is subtracted from the total weight of the thread and thrombus (see J.R. Smith, Br J Pharmacol, 5 77:29, 1982).

FeCl₃ model of arterial injury

The carotid arteries are isolated via a midline ventral cervical incision. A thermocouple is placed under each 10 artery and vessel temperature is recorded continuously on a strip chart recorder. A cuff of tubing (0.058 ID x 0.077 OD x 4 mm, Baxter Med. Grade Silicone), cut longitudinally, is placed around each carotid directly above the thermocouple. FeCl₃ hexahydrate is dissolved in water and the 15 concentration (20 percent) is expressed in terms of the actual weight of FeCl₃ only. To injure the artery and induce thrombosis, 2.85 µL is pipetted into the cuff to bathe the artery above the thermocouple probe. Arterial occlusion is indicated by a rapid drop in temperature. The 20 time to occlusion is reported in minutes and represents the elapsed time between application of FeCl₃ and the rapid drop in vessel temperature (see K.D. Kurz, Thromb. Res., 60:269, 1990).

25 Coagulation parameters

Plasma thrombin time (TT) and activated partial thromboplastin time (APTT) are measured with a fibrometer. Blood is sampled from a jugular catheter and collected in syringe containing sodium citrate (3.8 percent, 1 part to 9 30 parts blood). To measure TT, rat plasma (0.1 mL) is mixed with saline (0.1 mL) and bovine thrombin (0.1 mL, 30 U/mL in TRIS buffer; Parke Davis) at 37 °C. For APTT, plasma (0.1 mL) and APTT solution (0.1 mL, Organon Teknika) are incubated for 5 minutes (37 °C) and CaCl₂ (0.1 mL, 0.025 M) 35 is added to start coagulation. Assays are done in duplicate and averaged.

- 31 -

Index of Bioavailability

Bioavailability studies may be conducted as follows.

Compounds are administered as aqueous solutions to male
5 Fisher rats, intravenously (iv) at 5 mg/kg via tail vein
injection and orally (po) to fasted animals at 20 mg/kg by
gavage. Serial blood samples are obtained at 5, 30, 120,
and 240 minutes postdose following intravenous
administration and at 1, 2, 4, and 6 hours after oral
10 dosing. Plasma is analyzed for drug concentration using an
HPLC procedure involving C8 Bond Elute (Varion) cartridges
for sample preparation and a methanol/30 mM ammonium acetate
buffer (pH 4) gradient optimized for each compound. % Oral
bioavailability is calculated by the following equation:

15

$$\% \text{ Oral bioavailability} = \frac{\text{AUC po}}{\text{AUC iv}} \times \frac{\text{Dose iv}}{\text{Dose po}} \times 100$$

where AUC is area under the curve calculated from the plasma
level of compound over the time course of the experiment
20 following oral (AUC po) and intravenous (AUC iv) dosing.

Compounds

Compound solutions are prepared fresh daily in normal saline
and are injected as a bolus or are infused starting 15
25 minutes before and continuing throughout the experimental
perturbation which is 15 minutes in the arteriovenous shunt
model and 60 minutes in the FeCl₃ model of arterial injury
and in the spontaneous thrombolysis model. Bolus injection
volume is 1 mL/kg for i.v., and 5 mL/kg for p.o., and
30 infusion volume is 3 mL/hr.

Statistics

Results are expressed as means +/- SEM. One-way analysis of
variance is used to detect statistically significant
35 differences and then Dunnett's test is applied to determine

- 32 -

which means are different. Significance level for rejection of the null hypothesis of equal means is $P < 0.05$.

Animals

- 5 Male dogs (Beagles; 18 months - 2 years; 12-13 kg, Marshall Farms, North Rose, New York 14516) are fasted overnight and fed Purina certified Prescription Diet (Purina Mills, St. Louis, Missouri) 240 minutes after dosing. Water is available *ad libitum*. The room temperature is maintained
10 between 66-74 °F; 45-50 percent relative humidity; and lighted from 0600-1800 hours.

Pharmacokinetic model.

- Test compound is formulated immediately prior to dosing by
15 dissolving in sterile 0.9 percent saline to a 5 mg/mL preparation. Dogs are given a single 2 mg/kg dose of test compound by oral gavage. Blood samples (4.5 mL) are taken from the cephalic vein at 0.25, 0.5, 0.75, 1, 2, 3, 4 and 6 hours after dosing. Samples are collected in citrated
20 Vacutainer tubes and kept on ice prior to reduction to plasma by centrifugation. Plasma samples are analyzed by HPLC MS. Plasma concentration of test compound is recorded and used to calculate the pharmacokinetic parameters: elimination rate constant, K_e ; total clearance, Cl_t ; volume
25 of distribution, V_D ; time of maximum plasma test compound concentration, T_{max} ; maximum concentration of test compound of T_{max} , C_{max} ; plasma half-life, $t_{0.5}$; and area under the curve, A.U.C.; fraction of test compound absorbed, F .

30 Canine Model of Coronary Artery Thrombosis

- Surgical preparation and instrumentation of the dogs are as described in Jackson, et al., Circulation, 82, 930-940 (1990). Mixed-breed hounds (aged 6-7 months, either sex, Butler Farms, Clyde, New York, U.S.A.) are anesthetized with
35 sodium pentobarbital (30 mg/kg intravenously, i.v.), intubated, and ventilated with room air. Tidal volume and

- 33 -

respiratory rates are adjusted to maintain blood PO_2 , PCO_2 , and pH within normal limits. Subdermal needle electrodes are inserted for the recording of a lead II ECG.

- 5 The left jugular vein and common carotid artery are isolated through a left mediolateral neck incision. Arterial blood pressure (ABP) is measured continuously with a precalibrated Millar transducer (model MPC-500, Millar Instruments, Houston, TX, U.S.A.) inserted into the carotid artery. The
10 jugular vein is cannulated for blood sampling during the experiment. In addition, the femoral veins of both hindlegs are cannulated for administration of test compound.

- A left thoracotomy is performed at the fifth intercostal
15 space, and the heart is suspended in a pericardial cradle. A 1- to 2-cm segment of the left circumflex coronary artery (LCX) is isolated proximal to the first major diagonal ventricular branch. A 26-gauge needle-tipped wire anodal electrode (Teflon-coated, 30-gauge silverplated copper wire)
20 3-4 mm long is inserted into the LCX and placed in contact with the intimal surface of the artery (confirmed at the end of the experiment). The stimulating circuit is completed by placing the cathode in a subcutaneous (s.c.) site. An adjustable plastic occluder is placed around the LCX, over
25 the region of the electrode. A precalibrated electromagnetic flow probe (Carolina Medical Electronics, King, NC, U.S.A.) is placed around the LCX proximal to the anode for measurement of coronary blood flow (CBF). The occluder is adjusted to produce a 40-50 percent inhibition
30 of the hyperemic blood flow response observed after 10-s mechanical occlusion of the LCX. All hemodynamic and ECG measurements are recorded and analyzed with a data acquisition system (model M3000, Modular Instruments, Malvern, PA, U.S.A.).

35

- 34 -

Thrombus Formation and Compound Administration Regimens

Electrolytic injury of the intima of the LCX is produced by applying 100- μ A direct current (DC) to the anode. The current is maintained for 60 min and then discontinued whether the vessel has occluded or not. Thrombus formation proceeds spontaneously until the LCX is totally occluded (determined as zero CBF and an increase in the S-T segment). Compound administration is started after the occluding thrombus is allowed to age for 1 hour. A 2-hour infusion of the compounds of the present invention at doses of 0.5 and 1 mg/kg/hour is begun simultaneously with an infusion of thrombolytic agent (e.g. tissue plasminogen activator, streptokinase, APSAC). Reperfusion is followed for 3 hour after administration of test compound. Reocclusion of coronary arteries after successful thrombolysis is defined as zero CBF which persisted for at least 30 minutes.

Hematology and template bleeding time determinations

Whole blood cell counts, hemoglobin, and hematocrit values are determined on a 40- μ L sample of citrated (3.8 percent) blood (1 part citrate:9 parts blood) with a hematology analyzer (Cell-Dyn 900, Sequoia-Turner. Mount View, CA, U.S.A.). Gingival template bleeding times are determined with a Simplate II bleeding time device (Organon Teknika Durham, N.C., U.S.A.). The device is used to make 2 horizontal incisions in the gingiva of either the upper or lower left jaw of the dog. Each incision is 3 mm wide x 2 mm deep. The incisions are made, and a stopwatch is used to determine how long bleeding occurs. A cotton swab is used to soak up the blood as it oozes from the incision. Template bleeding time is the time from incision to stoppage of bleeding. Bleeding times are taken just before administration of test compound (0 min), 60 min into infusion, at conclusion of administration of the test compound (120 min), and at the end of the experiment.

- 35 -

All data are analyzed by one-way analysis of variance (ANOVA) followed by Student-Neuman-Kuels post hoc t test to determine the level of significance. Repeated-measures ANOVA are used to determine significant differences between
5 time points during the experiments. Values are determined to be statistically different at least at the level of $p < 0.05$. All values are mean \pm SEM. All studies are conducted in accordance with the guiding principles of the American Physiological Society. Further details regarding
10 the procedures are described in Jackson, et al., J. Cardiovasc. Pharmacol., (1993), 21, 587-599.

The following Examples are provided to further describe the invention and are not to be construed as
15 limitations thereof.

The abbreviations, symbols and terms used in the examples have the following meanings.

Ac = acetyl
AIBN = azobisisobutyronitrile
20 Anal. = elemental analysis
aq = aqueous
Bn or Bzl = benzyl
Boc = t-butyloxycarbonyl
Bu = butyl
25 n-BuLi = butyllithium
Calc = calculated
conc = concentrated
DCC = dicyclohexylcarbodiimide
DMAP = 4-dimethylaminopyridine
30 DMF = dimethylformamide
DMSO = dimethylsulfoxide
EDC = 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride
eq = (molar) equivalent
35 Et = ethyl
EtOAc = ethyl acetate

- 36 -

Et₃N = triethylamine
Et₂O = diethyl ether
EtOH = ethanol
Hex = hexanes
5 HOAt = 1-hydroxy-7-azabenzotriazole
HOBT = 1-hydroxybenzotriazole
HPLC = High Performance Liquid Chromatography
HRMS = high resolution mass spectrum
i-PrOH = isopropanol
10 IR = Infrared Spectrum
Me = methyl
MeI = methyl iodide
MeOH = methanol
MS(FAB) = fast atom bombardment mass spectrum
15 MS(FD) = field desorption mass spectrum
MS(IS) = ion spray mass spectrum
NBS = N-bromosuccinimide
NMR = Nuclear Magnetic Resonance
Ph = phenyl
20 i-Pr = isopropyl
RPHPLC = Reversed Phase High Performance Liquid
Chromatography
satd = saturated
SiO₂ = silica gel
25 TBS = tert-butyldimethylsilyl
TFA = trifluoroacetic acid
THF = tetrahydrofuran
TIPS = triisopropylsilyl
TLC = thin layer chromatography
30 tosyl = p-toluenesulfonyl
triflic acid = trifluoromethanesulfonic acid

Unless otherwise stated, pH adjustments and work up are
with aqueous acid or base solutions. ¹H-NMR indicates a
35 satisfactory NMR spectrum was obtained for the compound

- 37 -


described. IR indicates a satisfactory infra red spectrum was obtained for the compound described.

For consistency and clarity, a number of compounds are named as substituted diamine derivatives.

- 38 -

Example 1

Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine hydrochloride



5

A) 2-nitro-4-(dimethyl-t-butylsiloxy)aniline

To a mixture of 4-amino-3-nitrophenol (10.07 g, 65.3 mmol) and DMF (20 mL) was added imidazole (11.15 g, 163.8 mmol) followed by *t*-butyldimethylsilyl chloride (11.82 g, 78.4 mmol) in several portions. After 5 h, the reaction was diluted with EtOAc (150 mL) and washed with water (5 x 20 mL). The organic layer was MgSO₄, dried, filtered, and concentrated. The residue was chromatographed (10% EtOAc/hexanes to 20% EtOAc/hexanes) to give the title compound as a solid (17.06 g, 97%); mp 80-83 °C; IR (CHCl₃): 3399, 2932, 1519, 1242, 866 cm⁻¹; NMR (300 MHz, CDCl₃): δ 0.19 (s, 6H), 0.97 (s, 9H), 6.70 (d, 1H, J = 9.0), 6.95 (d, 1H, J = 3.0), 7.56 (d, 1H, J = 2.7); MS(FD): 268.2.

Analysis for C₁₂H₂₀N₂O₃Si:
Calculated: C 53.70, H 7.51, N 10.44
Found: C 53.47, H 7.50, N 10.31

B) N-(3-cyanobenzoyl)-2-nitro-4-(dimethyl-t-butylsiloxy)-aniline

- 39 -

- To a mixture of 2-nitro-4-(dimethyl-t-butylsiloxy)-aniline (10.01 g, 37.30 mmol) and CH₂Cl₂ (400 mL) at 0 °C was added pyridine (3.4 mL, 42.0 mmol) followed by the 3-cyanobenzoyl chloride (8.65 g, 52.2 mmol). The reaction was allowed to warm to room temperature and stirred for 1 h. The reaction was diluted with CH₂Cl₂ (800 mL) and washed with sat. aq. NH₄Cl (2 x 200 mL). The organic layer was MgSO₄ dried, filtered, and concentrated. The residue was chromatographed (20% EtOAc/hexanes) to give the title compound as a solid (13.94 g, 94%); IR(CHCl₃): 2932, 1689, 1508, 1289, 863 cm⁻¹; NMR(300 MHz, CDCl₃): δ 0.27 (s, 6H), 1.02 (s, 9H), 7.27 (s, 1H), 7.70 (m, 2H), 7.90 (d, J = 7.8, 1H), 8.16 (d, J = 1.2, 1H), 8.30 (s, 1H), 8.79 (d, J = 9.3, 1H), 11.15 (br s, 1H); MS(FD): 397.0.
- Analysis for C₂₀H₂₃N₃O₄Si:
Calculated: C 60.43, H 5.83, N 10.57
Found: C 60.58, H 5.79, N 10.61

- C) N¹-(3-cyanobenzoyl)-4-(dimethyl-t-butylsiloxy)-1,2-benzenediamine

A mixture of N-(3-cyanobenzoyl)-2-nitro-4-(dimethyl-t-butylsiloxy)aniline (5.00 g, 12.6 mmol), 10% Pd/C (2.5 g), EtOH (160 mL), and THF (160 mL) was hydrogenated at 1 atm for 12 h. The reaction was filtered through diatomaceous earth with hot methanolic washes. The filtrate was concentrated and chromatographed (20% EtOAc/hexanes to 30% EtOAc/hexanes) to give the title compound as a solid (2.72 g, 59%).

- D) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(dimethyl-t-butylsiloxy)-1,2-benzenediamine

To a mixture of 4-isopropyl benzoic acid (1.66 g, 10.1 mmol) and toluene (87 mL) was added pyridine (1.04 mL, 12.88 mmol) and thionyl chloride (0.96 mL, 13.15 mmol). The

- 40 -

reaction mixture was heated to 80 °C for 3 h, cooled, and concentrated. The crude 4-isopropyl benzoyl chloride was used without further purification as described below.

To a mixture of N¹-(3-cyanobenzoyl)-4-(dimethyl-t-butylsiloxy)-1,2-benzenediamine (2.63 g, 7.17 mmol) and CH₂Cl₂ (45 mL) at 0 °C was added pyridine (0.65 mL, 8.0 mmol) followed by a solution of the 4-isopropyl benzoyl chloride (10.1 mmol) in CH₂Cl₂ (90 mL). The reaction was allowed to warm to room temperature and stirred for 10 m. The reaction was diluted with CH₂Cl₂ (200 mL) and washed with sat. aq. NH₄Cl (2 x 50 mL). The organic layer was MgSO₄ dried filtered, and concentrated. The residue was chromatographed (10% EtOAc/hexanes to 20% EtOAc/hexanes) to give the title compound as a solid (3.04 g, 83%); IR(CHCl₃): 2962, 1656, 1610, 1513, 1473, 1295 cm⁻¹; NMR(300 MHz, CDCl₃): δ 0.16 (s, 6H), 0.95 (s, 9H), 1.29 (s, 3H), 1.31 (s, 3H), 3.00 (m, 1H), 6.68 (d, 1H, J = 9.0), 6.86 (s, 1H), 7.38 (d, 2H, J = 8.1), 7.50 (d, 1H, J = 9.0), 7.61 (m, 1H), 7.81 (d, 1H, J = 6.3), 7.89 (d, 2H, J = 7.1), 8.16 (d, 1H, J = 9.3), 8.29 (s, 1H), 8.58 (br s, 1H), 9.40 (br s, 1H); MS(FD): 513.1.

Analysis for C₃₀H₃₅N₃O₃Si:

Calculated: C 70.14, H 6.87, N 8.18

Found: C 70.30, H 7.01, N 8.35

25

E) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine

To a mixture of N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(dimethyl-t-butylsiloxy)-1,2-benzenediamine (1.57 g, 3.05 mmol) THF (30 mL) at 0 °C was added a THF solution of TBAF (3.1 mL, 3.1 mmol). After 1 h, the reaction was quenched with water (35 mL) and extracted with ether (2 x 200 mL). The combined organic layers was MgSO₄ dried, filtered, and concentrated. The residue was chromatographed

- 41 -

(1:1 EtOAc/hexanes) to give the title compound as a solid (1.10 g, 91%); IR(KBr): 3277, 1629, 1600, 1535, 1478, 1299 cm^{-1} ; NMR(300 MHz, d_6 -DMSO): δ 10.02 (s, 1H), 9.70 (s, 1H), 9.56 (s, 1H), 8.31 (s, 1H), 8.18 (d, $J = 7.5$, 1H), 8.01 (d, $J = 7.2$, 1H), 7.79 (d, $J = 8.1$, 2H), 7.70 (m, 1H), 7.34 - 7.21 (m, 4H), 6.63 (d, $J = 9.0$, 1H), 2.90 (m, 1H), 1.17 (s, 3H), 1.15 (s, 3H); MS(FD): 399.1.

Analysis for $\text{C}_{24}\text{H}_{21}\text{N}_3\text{O}_3$:

Calculated: C 72.17, H 5.30, N 10.52

10 Found: C 71.88, H 5.32, N 10.24

F) N^1 -[3-(amino(hydroxyimino)methyl)benzoyl]- N^2 -(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine

To a mixture of N^1 -(3-cyanobenzoyl)- N^2 -(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine (445 mg, 1.11 mmol) and absolute EtOH (30 mL) was added hydroxylamine hydrochloride (125 mg, 1.80 mmol) followed by triethylamine (0.3 mL, 2.15 mmol). The resulting mixture was heated to 80 °C for 2 h, cooled to room temperature, and filtered to give the title compound as a solid (238 mg, 0.55 mmol, 50%); NMR(300 MHz, d_6 -DMSO): δ 9.92 (s, 1H), 9.76 (s, 1H), 9.69 (s, 1H), 9.53 (s, 1H), 8.23 (s, 1H), 7.84 (m, 4H), 7.49 (m, 1H), 7.29 (m, 4H), 6.65 (d, $J = 9.0$, 1H), 5.85 (s, 2H), 2.92 (m, 1H), 1.19 (s, 3H), 1.17 (s, 3H); MS(FD): 432.2.

25 Analysis for $\text{C}_{24}\text{H}_{24}\text{N}_4\text{O}_4$:

Calculated: C 66.65, H 5.59, N 12.96

Found: C 65.75, H 5.43, N 12.42

G) N^1 -[3-(amino(imino)methyl)benzoyl]- N^2 -(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine hydrochloride

A mixture of N^1 -[3-(amino(hydroxyimino)methyl)benzoyl]- N^2 -(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine (209 mg, 0.48 mmol), EtOH (5 mL), H_2O (2.5 mL), 1N HCl (0.5 mL, 0.5 mmol), and 10% Pd/C (217 mg) was hydrogenated

- 42 -

at 1 atm. for 14 h. The reaction was filtered through diatomaceous earth with hot methanolic washes. The filtrate was concentrated and the residue was resubjected to the hydrogenation conditions using 10% Pd/C (189 mg). After 5 6 h, the reaction mixture was filtered through diatomaceous earth with hot methanolic washes. The filtrate was concentrated to give the title compound as a solid (134 mg, 62%); NMR(300 MHz, DMSO-d₆): δ 10.27 (s, 1H), 9.89 (s, 1H), 9.45 (s, 2H), 9.14 (s, 2H), 8.42 (s, 1H), 8.23 (d, J = 7.5, 10 1H), 7.94 (d, J = 7.2, 1H), 7.86 (d, J = 8.1, 2H), 7.73 (t, J = 7.8, 1H), 7.28 (m, 4H), 6.65 (d, J = 8.7, 1H), 2.90 (m, 1H), 1.19 (s, 3H), 1.17 (s, 3H); MS(FD): 399.2 (M-OH).


Analysis for C₂₄H₂₄N₄O₃ · HCl · 2.5 H₂O :

C, 57.89, H, 6.07, N, 11.25

15 Found: C, 57.21, H, 5.32, N, 10.90

Example 2

Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-carboxymethoxy-1,2-benzenediamine
20 dihydrochloride



A) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-
25 (benzyloxycarbonylmethoxy)-1,2-benzenediamine

- 43 -

To a mixture of N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine (502 mg, 1.26 mmol), acetone (15 mL), and potassium carbonate (216 mg, 1.56 mmol) was added by benzyl 2-bromoacetate (0.4 mL, 2.52 mmol). After stirring overnight, the reaction was concentrated and the residue was chromatographed (100% CH₂Cl₂ to 10% EtOAc/CH₂Cl₂) to give the title compound as a solid (532 mg, 77%); IR(CHCl₃): 2965, 1759, 1654, 1610, 1514, 1175 cm⁻¹; NMR(300 MHz, CDCl₃): δ 1.28 (s, 3H), 1.30 (s, 3H), 3.00 (m, 1H), 4.50 (s, 2H), 5.22 (s, 2H), 6.71 (d, 1H, J = 9.3), 6.93 (s, 1H), 7.31 (m, 7H), 7.54 (d, 1H, J = 21), 7.60 (t, 1H, J = 7.8), 7.78 (d, 1H, J = 8.7), 7.89 (d, 2H, J = 8.1), 8.19 (d, 1H, J = 8.1), 8.30 (s, 1H), 8.74 (br s, 1H), 9.48 (br s, 1H); MS(FD): 547.

15 Analysis for C₃₃H₂₉N₃O₅:

Calculated: C, 72.38, H, 5.34, N, 7.67

Found: C, 72.55, H, 5.32, N, 7.78

20 B) N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(benzyloxycarbonylmethoxy)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting with N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(benzyloxycarbonylmethoxy)-1,2-benzenediamine (532 mg, 0.97 mmol), a residue was obtained after concentrating the reaction mixture. The residue was chromatographed (10% EtOAc/CH₂Cl₂ to 50% EtOAc/CH₂Cl₂) to give the title compound as a solid (317 mg, 56%); NMR(300 MHz, DMSO-d₆): δ 1.17 (s, 3H), 1.19 (s, 3H), 2.92 (m, 1H), 4.88 (s, 2H), 5.20 (s, 2H), 5.89 (br s, 2H), 6.85 (d, 1H, J = 9.0), 7.4 (m, 11H), 7.83 (d, 2H, J = 7.5), 7.88 (d, 1H, J = 8.1), 8.24 (s, 1H), 9.73 (s, 1H), 9.88 (s, 1H), 10.01 (s, 1H); MS(FD): 580.0.

Analysis for C₃₃H₃₂N₄O₆ · 0.35 H₂O:

Calculated for: C, 67.53, H, 5.62, N, 9.55

- 44 -

Found: C, 68.13, H, 5.69, N, 9.49

C) N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-carboxymethoxy-1,2-benzenediamine dihydrochloride

5 Using a procedure similar to Example 1G except starting
with N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-
isopropylbenzoyl)-4-(benzyloxy-carbonylmethoxy)-1,2-
benzenediamine, the title compound was obtained as a solid
(62%); NMR(300 MHz, d₆-DMSO): δ 1.17 (s, 3H), 1.19 (s, 3H),
10 2.92 (m, 1H), 4.68 (s, 2H), 6.84 (d, 2H, J = 9.0 Hz), 7.34
(d, 2H, J = 8.4 Hz), 7.44 (d, 1H, J = 8.7 Hz), 7.74 (t, 1H,
J = 7.8 Hz), 7.89 (d, 2H, J = 7.8 Hz), 7.96 (d, 1H, J = 8.4
Hz), 8.26 (d, 1H J = 7.8 Hz), 8.42 (s, 1H), 9.15 (s, 2H),
9.45 (s, 2H), 10.01 (s, 1H), 10.36 (s, 1H); MS(FD): 474.


15 Analysis for $C_{26}H_{26}N_4O_5 \cdot 2 HCl \cdot H_2O$:

Calculated: C, 55.23, H, 5.35, N, 9.91

Found: C, 55.36, H, 5.07, N, 9.90

Example 3

20 Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-
(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-
benzenediamine hydrochloride



25

- 45 -

A) N^1 -(3-cyanobenzoyl)- N^2 -(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine

Using a procedure similar to Example 2A except starting with N^1 -(3-cyanobenzoyl)- N^2 -(4-isopropylbenzoyl)-4-hydroxy-1,2-benzenediamine and ethyl bromoacetate, the title compound was obtained as a solid (78%); IR(CHCl₃): 2966, 1757, 1655, 1610, 1514 cm⁻¹; NMR(300 MHz, CDCl₃): δ 1.31 (m, 9H), 3.00 (m, 1H), 4.24 (t, 2H, J = 6.9 Hz), 4.35 (s, 2H), 6.59 (d, 1H, J = 9.0 Hz), 6.95 (s, 1H), 7.35 (m, 3H), 7.64 (m, 1H), 7.82 (d, 1H, J = 7.5 Hz), 7.92 (d, 2H, J = 8.1 Hz), 8.23 (d, 1H, J = 7.5 Hz), 8.32 (s, 1H), 9.06 (br s, 1H), 9.61 (br s, 1H); MS(FD): 485.2.

Analysis for C₂₈H₂₇N₃O₅:

Calculated: C, 69.26, H, 5.60, N, 8.65

Found: C, 69.52, H, 5.71, N, 8.58

B) N^1 -[3-(amino(hydroxyimino)methyl)benzoyl]- N^2 -(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting with N^1 -(3-cyanobenzoyl)- N^2 -(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine (632 mg, 1.30 mmol), the title compound was obtained as a solid (60%); IR(KBr): 3345, 1753, 1647, 1521, 1178 cm⁻¹; NMR(300 MHz, DMSO-d): δ 1.19 (m, 9H), 2.89 (m, 1H), 4.17 (q, 2H, J = 7.2 Hz), 4.78 (s, 2H), 5.88 (s, 2H), 6.83 (d, 1H, J = 9.0 Hz), 7.49-7.34 (m, 6H), 7.81 (d, 2H, J = 8.1 Hz), 7.86 (d, 1H, J = 12.9 Hz), 8.24 (s, 1H), 9.72 (s, 1H), 9.86 (s, 1H), 10.00 (s, 1H); MS(FD): 518.1.

Analysis for C₂₈H₃₀N₄O₆:

Calculated: C, 64.85, H, 5.83, N, 10.80

Found: C, 64.59, H, 5.80, N, 10.59

- 46 -

C) N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 1G except starting with N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine, the title compound was obtained as a solid (65%); NMR(300 MHz, d₆-DMSO): δ 1.03 (t, 3H, J = 7.1 Hz), 1.17 (s, 3H), 1.19 (s, 3H), 2.92 (m, 1H), 4.17 (q, 2H, J = 6.9 Hz), 4.78 (s, 2H), 6.83 (d, 1H, J = 9.0 Hz), 7.34 (d, 3H, J = 7.8 Hz), 7.45 (d, 1H, J = 8.7 Hz), 7.74 (t, 1H, J = 7.8 Hz), 7.88 (d, 2H, J = 7.8 Hz), 7.95 (d, 1H, J = 8.4 Hz), 8.24 (d, 1H, J = 7.8 Hz), 8.42 (s, 1H), 9.13 (s, 2H), 9.45 (s, 2H), 9.99 (s, 1H), 10.34 (s, 1H); MS(FD): 503.


Analysis for C₂₈H₃₀N₄O₅ · HCl · H₂O:

Calculated: C, 60.38, H, 5.97, N, 10.06

Found: C, 60.64, H, 5.72, N, 9.97

Example 4

Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride



A) N¹-[3-cyanobenzoyl]-2-nitroaniline

Using a procedure similar to Example 1B except starting with 2-nitroaniline, the title compound was obtained as a yellow solid (98%), mp 154-155.5 °C; NMR (DMSO): δ 7.47 (dt, 1H, J=9.0, 2.4 Hz), 7.69-7.83 (m, 3H), 8.03 (dd, 1H,

- 47 -

J=9.9, 1.8 Hz), 8.13 (dt, J=9.3, 1.2 Hz), 8.25 (dt, 1H, J=9.9, 1.5 Hz), 8.39 (s, 1H), 10.92 (s, 1H).

Analysis for C₁₄H₉N₃O:

Calculated: C, 62.92; H, 3.39; N, 15.72

5 Found: C, 62.86; H, 3.44; N, 16.02

B) N¹-[3-cyanobenzoyl]-1,2-benzenediamine

Using a procedure similar to Example 1C except starting with N¹-[3-cyanobenzoyl]-2-nitroaniline, the title compound
10 was obtained as a yellow solid (72%), mp 197-200 °C; NMR (DMSO): δ 5.00 (s, 2H), 6.59 (t, 1H, J=9.3 Hz), 6.79 (d, 1H, J=9.4 Hz), 6.98 (t, 1H, J=9.3 Hz), 7.17 (d, 1H, J=9.4 Hz), 7.73 (t, 1H, J=10.2 Hz), 8.05 (d, 1H, J=10.1 Hz), 8.27 (d, 1H, J=10.1 Hz), 8.44 (s, 1H), 9.81 (s, 1H); MS(FD): 237.

15 Analysis for C₁₄H₁₁N₃O:

Calculated: C, 70.87; H, 4.67; N, 17.71

Found: C, 70.68; H, 4.58; N, 17.52

C) N¹-[3-cyanobenzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine
20

Using a procedure similar to Example 1B except starting with N¹-[3-cyanobenzoyl]-1,2-benzenediamine and using triethylamine as base, the title compound was obtained as a white solid (730 mg, 87%); mp 174-174.5 °C; NMR (DMSO): δ
25 1.27 (s, 9H), 7.27-7.32 (m, 2H), 7.53 (d, 2H, J=10.2 Hz), 7.61-7.64 (m, 1H), 7.86-7.72 (m, 1H), 7.75 (t, 1H, J=9.3 Hz), 7.88 (d, 2H, J=10.2 Hz), 8.06 (dd, 1H, J=7.8, 1.5 Hz), 8.23 (dd, 1H, J=9.6, 1.8 Hz), 8.36 (s, 1H), 9.90 (br s, 1H), 10.24 (br s, 1H); MS(FD+):

30 Analysis for C₂₅H₂₃N₃O₂:

Calculated: C, 75.54; H, 5.83; N, 10.57

Found: C, 75.69; H, 6.14; N, 10.57

- 48 -

D) N^1 -[3-(amino(hydroxyimino)methyl)benzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting
5 N^1 -[3-cyanobenzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a white solid (55%), mp 201-204 °C; NMR(DMSO): δ 1.30 (s, 9H), 5.91 (br s, 2H), 7.28-7.32 (m, 2H), 7.50-7.56 (m, 3H), 7.61-7.67 (m, 1H), 7.68-7.73 (m, 1H), 7.86-7.94 (m, 4H), 8.28 (s, 1H),
10 9.75 (s, 1H), 9.96 (s, 1H), 10.14 (s, 1H); MS(FD+): .

Analysis for $C_{25}H_{26}N_4O_3$:

Calculated: C, 69.75; H, 6.09; N 13.01,

Found: C, 69.48; H, 5.91; N, 12.74

15

E) N^1 -[3-(amino(imino)methyl)benzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 1G except starting
 N^1 -[3-(amino(hydroxyimino)methyl)benzoyl]- N^2 -(4-t-butyl-
20 benzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (76%); mp 196-198 °C; NMR (DMSO): δ 1.32 (s, 9H), 7.28-7.32 (m, 2H), 7.52 (d, 2H, J=10.2 Hz), 7.62-7.67 (m, 1H), 7.67-7.71 (m, 1H), 7.77 (t, 1H, J=9.3 Hz), 7.96 (d, 2H, J=10.2 Hz), 8.00 (d, 1H, J=10.8 Hz), 8.30 (d, 1H, J=9.3
25 Hz), 8.49 (s, 1H), 9.20 (s, 1H), 9.49 (s, 1H), 10.13 (s, 1H), 10.56 (s, 1H); MS(FD+): 415.

Analysis for $C_{25}H_{26}N_4O_2 \cdot HCl \cdot 0.75H_2O$:

Calculated C, 64.03; H, 6.02; N, 11.95

Found: C, 64.71; H, 6.19; N, 12.16

30

Example 5

Preparation of N^1, N^2 -bis[3-(aminoiminomethyl)benzoyl]-1,2-benzenediamine dihydrochloride

- 49 -

A) N¹,N²-bis(3-cyanobenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1D, except starting with 1,2-benzenediamine and 3-cyanobenzoyl chloride with triethylamine as base, the title compound was obtained as an off-white solid (18%); IR(KBr): 3362, 2238, 1656, 1552 cm⁻¹; NMR(300 MHz, DMSO-d₆): 7.33 (m, 2H); 7.64 (m, 2H); 7.73 (t, 2H, J=9.5 Hz); 8.05 (d, 2H, J=9.3 Hz); 8.24 (d, 2H, J=9.9 Hz); 8.35 (s, 2H); 10.18 (br s, 2H); MS(FD): 366.1.

Analysis for C₂₂H₁₄N₄O₂:

Calculated: C, 72.13; H, 3.85; N, 15.29
Found: C, 71.86; H, 3.96; N, 15.35

B) N¹,N²-bis[3-(amino(hydroxyiminomethyl))benzoyl]-1,2-benzenediamine

Using a procedure similar to Example 1F except starting with N¹,N²-bis(3-cyanobenzoyl)-1,2-benzenediamine, the title compound was obtained as a white solid (92%); IR(KBr): 3324, 1658, 1514 cm⁻¹; NMR(300 MHz, DMSO-d₆): 5.89 (br s, 4H); 7.31 (m, 2H); 7.52 (t, 2H, J=9.6 Hz); 7.68 (m, 2H); 7.85 (d, 2H, J=9.6 Hz); 7.93 (d, 2H, J=9.6 Hz); 8.27 (s, 2H); 9.74 (s, 2H); 10.08 (s, 2H); MS(FD): 432.1.

Analysis for C₂₂H₂₀N₆O₄:

Calculated: C, 61.10; H, 4.66; N, 19.43
Found: C, 61.25; H, 4.81; N, 19.14

C) N¹,N²-bis[3-(aminoiminomethyl)benzoyl]-1,2-benzenediamine dihydrochloride

Using a procedure similar to Example 1G except starting with N¹,N²-bis[3-(amino(hydroxyimino)methyl)benzoyl]-1,2-benzenediamine, the title compound was obtained as a tan solid (54%); IR(KBr): 1444, 1492, 1520, 1658 cm⁻¹; NMR(300 MHz, DMSO-d₆): 7.34 (m, 2H); 7.75 (m, 3H); 8.01 (d, 2H, J=9.0 Hz); 8.35 (d, 2H, J=8.7 Hz); 8.57 (s, 4H); 9.27 (s, 4H); 9.51 (s, 4H); 10.64 (s, 2H); MS(FD): 400.

- 50 -

Analysis for $C_{22}H_{26}N_6O_2 \cdot 2HCl$:

Calculated: C, 55.82; H, 4.68; N, 17.75

Found: C, 48.36; H, 4.69; N, 14.98

5

Example 6

Preparation of N^1 -[3-(amino(imino)methyl)benzoyl]- N^2 -(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine hydrochloride

- 10 A) N-(3-cyanobenzoyl)-4-(ethoxycarbonylmethyl)-2-nitroaniline

Using a procedure similar to Example 1B except starting with 4-(ethoxycarbonylmethyl)-2-nitroaniline, the title compound was obtained as a solid (98%); IR($CHCl_3$): 1340, 1516, 1694, 1733 cm^{-1} ; NMR(300 MHz, $CDCl_3$): 1.28 (t, 3H, J=7.2 Hz); 3.70 (s, 2H); 4.19 (q, 2H, J=7.2 Hz); 7.69 (m, 2H); 7.89 (d, 1H, J=8.1 Hz); 8.18 (d, 1H, J=7.8 Hz); 8.24 (s, 1H); 8.30 (s, 1H); 8.92 (d, 1H, J=9.3 Hz); MS(FD): 353.

15

Analysis for $C_{18}H_{15}N_3O_5$:

20

Calculated: C, 61.19; H, 4.33; N, 11.80

Found: C, 61.28; H, 4.35; N, 11.79

- B) N^1 -(3-cyanobenzoyl)-4-(ethoxycarbonylmethyl)-1,2-benzenediamine

25

Using a procedure similar to Example 1C except starting with N-(3-cyanobenzoyl)-4-(ethoxycarbonylmethyl)-2-nitroaniline, the title compound was obtained as a solid (78%); IR($CHCl_3$): 1519, 1673, 1728, 2235, 3025 cm^{-1} ; NMR(300 MHz, $DMSO-d_6$): 1.19 (t, 3H, J=8.4 Hz); 3.50 (s, 2H); 4.08 (q, 2H, J=8.4 Hz); 5.04 (s, 2H); 6.47 (d, 1H, J=9.6 Hz); 6.67 (s, 1H); 7.08 (d, 1H, J=9.6 Hz); 7.73 (t, 1H, J=9.6 Hz); 8.04 (d, 1H, J=9.6 Hz); 8.26 (d, 1H, J=9.9 Hz); 8.44 (s, 1H); 9.77 (s, 1H); MS(FD): 323.0.

30

Analysis for $C_{18}H_{17}N_3O_3$:

- 51 -

Calculated: C, 66.86; H, 5.30; N, 13.00

Found: C, 68.87; H, 5.70; N, 12.75

- 5 C) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(ethoxycarbonylmethyl)-1,2-benzenediamine

Using a procedure similar to Example 1D except starting with N-(3-cyanobenzoyl)-4-(ethoxycarbonylmethyl)-1,2-benzenediamine, the title compound was obtained as a solid (88%); IR(CHCl₃): 1308, 1524, 1654, 1729, 2966 cm⁻¹;
10 NMR(300 MHz, DMSO-d₆): 1.21 (m, 9H); 2.92 (m, 1H); 3.72 (s, 2H); 4.12 (q, 2H, J=6.9 Hz); 7.19 (d, 1H, J=8.4 Hz); 7.39 (d, 2H, J=8.4 Hz); 7.58 (d, 1H, J=8.4 Hz); 7.64 (s, 1H); 7.75 (t, 1H, J=7.5 Hz); 7.89 (d, 2H, J=8.4 Hz); 8.07 (d, 1H, J=8.1 Hz); 8.24 (d, 1H, J=7.8 Hz); 8.36 (s, 1H); MS(FD):
15 469.

Analysis for C₂₈H₂₇N₃O₄:

Calculated: C, 71.63; H, 5.80; N, 8.95

Found: C, 71.69; H, 6.02; N, 8.75

- 20 D) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine

To a mixture of N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(ethoxycarbonylmethyl)-1,2-benzenediamine (1.41 g, 3.00 mmol) in THF (30 mL) and MeOH (10 mL) was added 1 M
25 LiOH (10 mL) and stirred for 2.5 h. The reaction was diluted with Et₂O (300 mL) and washed with 1N HCl (40 mL) and H₂O (40 mL). The organic layer was concentrated to give the title compound as a pale yellow solid (1.32 g, quant.);
IR(KBr): 1326, 1515, 1625, 1672, 3326 cm⁻¹; NMR(300 MHz,
30 DMSO-d₆): 1.19 (d, 6H, J=6.9 Hz); 2.97 (m, 1H); 3.65 (s, 2H); 7.12 (d, 1H, J=8.1 Hz); 7.35 (d, 2H, J=7.8 Hz); 7.49 (d, 1H, J=8.1 Hz); 7.54 (s, 1H); 7.71 (t, 1H, J=8.0 Hz); 7.86 (d, 2H, J=7.8 Hz); 8.02 (d, 1H, J=7.2 Hz); 8.22 (d, 1H,

- 52 -

J=8.4 Hz); 8.34 (s, 1H); 9.96 (s, 1H); 10.27 (s, 1H);

MS(FD): 441.0.

Analysis for C₂₆H₂₃N₃O₄·1.5 H₂O:

Calculated: C, 67.30; H, 5.54; N, 9.06

5 Found: C, 67.31; H, 5.20; N, 8.92

E) N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting
10 with N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine, the title compound was obtained as a solid (21%); NMR(300 MHz, DMSO-d₆): 1.17 (m, 9H); 2.95 (m, 1H); 3.69 (s, 2H); 4.08 (q, 2H, J=7.2 Hz); 5.89 (s, 2H); 7.17 (d, 1H, J=8.4 Hz); 7.36 (d, 2H, J=8.1 Hz); 7.53 (m,
15 3H); 7.83 (m, 4H); 8.24 (s, 1H); 9.73 (s, 1H); 9.94 (s, 1H); 10.07 (s, 1H); MS(FD): 502.

Analysis for C₂₈H₃₀N₄O₅:

Calculated: C, 66.92; H, 6.02; N, 11.15

Found: C, 64.70; H, 5.74; N, 10.25

20

F) N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 1G except starting
with N¹-[3-(amino(hydroxy-imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-4-(carboxymethyl)-1,2-benzenediamine, the
25 title compound was obtained as a pale yellow solid (81%); NMR(300 MHz, DMSO-d₆): 1.16 (m, 9H); 2.91 (m, 1H); 3.68 (s, 2H); 4.07 (q, 2H, J=7.2 Hz); 7.15 (d, 1H, J=8.1 Hz); 7.33 (d, 2H, J=8.4 Hz); 7.57 (m, 3H); 7.73 (t, 1H, J=7.8 Hz);
30 7.91 (d, 2H, J=7.8 Hz); 7.96 (d, 1H, J=8.1 Hz); 8.26 (d, 1H, J=7.5 Hz); 8.45 (s, 1H); 9.19 (s, 2H); 9.47 (s, 2H); 10.10 (s, 1H); 10.52 (s, 1H); MS(FD): 487.0.

Analysis for C₂₈H₃₀N₄O₄·1.5 HCl·0.25 H₂O:

Calculated: C, 61.62; H, 5.91; N, 10.27

- 53 -

Found: C, 61.47; H, 5.58; N, 10.37

Example 7**Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-1,2-benzenediamine hydrochloride****A) N¹-(t-butoxycarbonyl)-1,2-benzenediamine**

To a mixture of BOC₂O (24.3 g, 111.2 mmol), 1:1 THF: H₂O (500 mL) and potassium carbonate was added phenylene diamine (10 g, 92.5 mmol) and stirred for 24 h. The reaction was concentrated and extracted with EtOAc. The organic layer was MgSO₄ dried and concentrated. The residue was chromatographed (10% EtOAc/hexanes to 40% EtOAc/hexanes) to give the title compound as a solid (14.56 g, 76%).

B) N¹-(t-butoxycarbonyl)-N²-(4-isopropylbenzoyl)-1,2-benzenediamine

Using a procedure similar to 1D except starting with N¹-(t-butoxycarbonyl)-1,2-benzenediamine, the title compound was obtained as a solid (85%); NMR(300 MHz, DMSO-d₆): 1.24 (d, 6H, J=8.4 Hz), 1.44 (s, 9H), 2.98 (m, 1H), 7.17 (m, 2H), 7.41 (d, 2H, J=9.9 Hz), 7.54 (d, 2H, J=9.0 Hz), 7.89 (d, 2H, J=9.6 Hz), 8.65 (s, 1H), 9.77 (s, 1H); MS(FD⁺): 354.2

25 Analysis for C₂₁H₂₆N₂O₃:

Calculated: C, 71.16; H, 7.39 N, 7.90

Found: C, 72.17; H, 7.53; N, 8.17

C) N¹-(4-isopropylbenzoyl)-1,2-benzenediamine

Using a procedure similar to 8K except starting with N¹-(t-butoxycarbonyl)-1,2-benzenediamine, the title compound was obtained as a solid (70%); NMR(300 MHz, CDCl₃): 1.29 (d, 6H, J=8.4 Hz), 2.99 (m, 1H), 6.81-6.87 (m, 2H), 7.10 (d,

- 54 -

1H, J=9.3 Hz), 7.34 (m, 3H), 7.84 (d, 2H, J=9.9 Hz);

MS(FD⁺): 254.

Analysis for C₁₆H₁₈N₂O:

Calculated: C, 75.56; H, 7.13 N, 11.01

5 Found: C, 75.81; H, 6.86; N, 10.75

D) N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-1,2-benzenediamine

Using a procedure similar to 1B except starting with
10 N¹-(4-isopropylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (53%); NMR(300 MHz, CDCl₃): 1.30 (d, 6H, J=8.4 Hz), 3.00 (m, 1H), 7.04 ((m, 2H), 7.34 (d, 1H, J=8.4 Hz), 7.38 (d, 1H, J=9.9 Hz), 7.55 (d, 1H, J=8.4 Hz), 7.62 (d, 1H, J=9.3 Hz), 7.83 (d, 1H, J=9.3 Hz),
15 7.91 (d, 2H, J=9.9 Hz), 8.23 (d, 1H, J=9.6 Hz), 8.31 (d, 2H, J=9.0 Hz), 8.83 (s, 1H), 9.78 (s, 1H); MS(FD⁺):383.3.

Analysis for C₂₄H₂₁N₃O₂·0.25 H₂O:

Calculated: C, 74.30; H, 5.59 N, 10.83

Found: C, 74.16; H, 5.55; N, 10.79

20

E) N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-1,2-benzenediamine

Using a procedure similar to 1F except starting with
N¹-(3-cyanobenzoyl)-N²-(4-isopropylbenzoyl)-1,2-benzene-
25 diamine, the title compound was obtained as a solid (40%);

Analysis for C₂₄H₂₁N₃O₂·0.25 H₂O:

Calculated: C, 74.30; H, 5.59 N, 10.83

Found: C, 74.16; H, 5.55; N, 10.79

30 F) N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-isopropylbenzoyl)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 1G except starting with N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-

- 55 -

isopropylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (63%); mp 227-9.0; MS(FD): 401.1.

Analysis for $C_{24}H_{25}N_4O_2 \cdot HCl \cdot 1.5 H_2O$:

Calculated: C, 62.13; H, 6.08; N, 12.08

5 Found: C, 61.90; H, 5.72; N, 11.91

Example 8

Preparation of N^1 -[(1-iminoisoindolin-6-yl)carbonyl]- N^2 -(*t*-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine trifluoroacetate

10

A) methyl 3-bromo-4-methylbenzoate

To a mixture of hexane washed 60% disp. NaH (2.28 g, 57 mmol) in DMF (250 mL), 0 °C, was added 3-bromo-4-methylbenzoic acid (10.03 g, 46.6 mmol). After 30 min, the reaction was warmed to room temperature, diluted with EtOAc (1.25 L) and washed with H₂O (5 x 250 mL). The organic layer was MgSO₄ dried and concentrated. The residue was chromatographed (~375 g silica, 10% EtOAc/CH₂Cl₂) to give the desired product as a solid (16.68 g, 63%); IR(CHCl₃): 1260, 1294, 1437, 1721 cm⁻¹; NMR(300 MHz, CDCl₃): 2.46 (s, 3H); 3.92 (s, 3H); 7.31 (d, 1H, J=7.8 Hz); 7.88 (d, 1H, J=7.8 Hz); 8.21 (s, 1H); MS(FD): 228.0.

15

20

Analysis for C₉H₉BrO₂:

25 Calculated: C, 47.19; H, 3.96

Found: C, 47.06; H, 3.92

B) methyl 3-cyano-4-methylbenzoate

A mixture of copper (I) cyanide (510 mg, 5.69 mmol) and DMF (16 mL) was heated to 140 °C. The reaction was added methyl 3-bromo-4-methylbenzoate (505 mg, 2.2 mmol) in DMF (10 mL) and heating was continued for 12 h. The reaction was cooled, diluted with EtOAc (125 mL) and washed with H₂O (5 x 25 mL). The EtOAc layer was dried over MgSO₄ and

30

- 56 -

concentrated. The residue was chromatographed (40 g silica, 10% EtOAc/hexanes) to give the title compound as a solid (236 mg, 61%); IR(CHCl₃): 1128, 1269, 1300, 1725, 2231, 3024 cm⁻¹; NMR(300 MHz, CDCl₃): 2.63 (s, 3H); 3.95 (s, 3H); 5 7.43 (d, 1H, J=8.1 Hz); 8.14 (d, 1H, J=8.1 Hz); 8.28 (s, 1H); MS(FD): 175.1.

Analysis for C₁₀H₉N₁O₂:

Calculated: C, 68.56; H, 5.18; N, 8.00

Found: C, 68.29; H, 5.18; N, 8.04

10

C) methyl 3-cyano-4-(bromomethyl)benzoate

A mixture of methyl 3-cyano-4-methylbenzoate (1.78 g, 9.79 mmol) and NBS (1.92 g, 10.8 mmol) in CCl₄ (150 mL) was irradiated with a 150W tungsten lamp at 50 C for 2 h. The 15 reaction was cooled, filtered, and concentrated. The residue was chromatographed (250 g silica, 10% EtOAc/hexanes to 15% EtOAc/hexanes) to give the title compound (1.20 g, 48%); IR(CHCl₃): 1291, 1302, 1439, 1728 cm⁻¹; NMR(300 MHz, CDCl₃): 4.01 (s, 3H); 4.67 (s, 3H); 7.66 (d, 20 1H, J=8.1 Hz); 8.24 (d, 1H, J=8.7 Hz); 8.34 (s, 1H); MS(FD): 253.0.

Analysis for C₁₀H₈NO₂:

Calculated: C, 47.27; H, 3.17; N, 5.51

Found: C, 47.45; H, 3.38; N, 5.53

25

D) methyl 3-cyano-4-[(N,N-bis(t-butoxycarbonyl)amino)-methyl]benzoate

To a mixture of hexane washed 60% disp. NaH (40 mg, 1 mmol) and DMF (8 mL) was added HN(BOC)₂ (203 mg, 0.93 mmol) and stirred for 30 m.. The reaction was added methyl 30 3-cyano-4-(bromomethyl)benzoate (202 mg, 0.80 mmol). After 3.5 h, the reaction was diluted with EtOAc (40 mL) and washed with H₂O (6 x 5 mL). The organic layer was MgSO₄ dried and concentrated. The crude material was

- 57 -

chromatographed (30 g silica, 10% EtOAc/hexanes) to give the title compound (quant.). IR(CHCl₃): 1125, 1370, 1723, 1791, 3020 cm⁻¹; NMR(300 MHz, CDCl₃): 1.46 (s, 18H); 3.96 (s, 3H); 5.08 (s, 2H); 7.42 (d, 1H, J=8.4 Hz); 8.23 (d, 1H, J=8.4 Hz); 8.32 (s, 1H); MS(FD): 391.

Analysis for C₂₀H₂₆N₂O₆:

Calculated: C, 61.53; H, 6.71; N, 7.17

Found: C, 61.26; H, 6.99; N, 7.13

- 10 E) 3-cyano-4-(N,N-bis(t-butoxycarbonyl)aminomethyl)benzoic acid

To a -25 °C solution of methyl 3-cyano-4-(N,N-bis-(t-butoxycarbonyl)aminomethyl)benzoate (1.81 g, 4.64 mmol) in THF (14 mL) was added a 1:1 mixture of MeOH (4.6 mL) and 15 1 M LiOH (4.6 mL). After 1.25 h, the reaction was quenched with 1 N HCl (4.6 mL), diluted with Et₂O (150 mL) and washed with H₂O (2 x 30 mL). The combined aqueous layers was extracted with EtOAc (150 mL). The combined organic layers was MgSO₄ dried, filtered, and concentrated. The crude 20 material was chromatographed (0.5% HOAc in 30% EtOAc/hexanes to 0.5% HOAc in 40% EtOAc/hexanes) to give the title compound (767 mg, 44%); IR(CHCl₃): 1146, 1370, 1704, 2984, 3020 cm⁻¹; NMR(300 MHz, DMSO-d₆): 1.36 (s, 18H); 4.93 (s, 2H); 7.41 (d, 1H, J=8.4 Hz); 8.21 (d, 1H, J=8.4 Hz); 8.25 25 (s, 1H); MS(FD): 377.0.

Analysis for C₁₉H₂₄N₂O₆:

Calculated: C, 60.63; H, 6.43; N, 7.44

Found: C, 60.39; H, 6.30; N, 7.36

- 30 F) N-(t-butoxycarbonyl)-2-nitro-4-(dimethyl-t-butyl-siloxy)aniline

To a mixture 2-nitro-4-(dimethyl-t-butylsiloxy)aniline (10.00 g, 37.3 mmol) in THF (85 mL) was added a solution of 1M NaHMDS/THF (75 mL, 75 mmol). After 15 minutes, a

- 58 -

solution of di-*t*-butyl dicarbonate (7.33 mg, 33.6 mmol) in THF (120 mL) was added and stirred for 30 m. The reaction was concentrated, diluted with EtOAc (350 mL) and washed with 0.1N HCl (850 mL). The aqueous layer was added sat. aq. NaHCO₃ (85mL) and extracted with EtOAc (2 x 350 mL). The combined EtOAc layers was MgSO₄ dried, filtered, and concentrated. The crude material chromatographed(hexanes to 5% EtOAc/ hexanes) the title compound (7.99 g, 21.7 mmol, 58%). This material was used without further purification.

10

G) N¹-(*t*-butoxycarbonyl)-4-(dimethyl-*t*-butylsiloxy)-1,2-benzenediamine

Using a procedure similar to Example 1C except starting with N-(*t*-butoxycarbonyl)-2-nitro-4-(dimethyl-*t*-butylsiloxy)aniline, the title compound was obtained (48%); IR(CHCl₃): 866, 1160, 1515, 1709, 2958 cm⁻¹; NMR(300 MHz, CDCl₃): 0.15 (s, 6H); 0.95 (s, 9H); 1.51 (s, 9H); 6.02 (br s, 2H); 6.24 (d, 1H, J=8.3 Hz); 6.27 (s, 1H); 6.99 (d, 1H, J=8.3 Hz); MS(FD): 338.3.

20 Analysis for C₁₇H₃₀N₂O₃:

Calculated: C, 60.32; H, 8.93; N, 8.28

Found: C, 60.16; H, 9.01; N, 8.26

25 H) N¹-(*t*-butoxycarbonyl)-N²-(*t*-butylbenzoyl)-4-(dimethyl-*t*-butylsiloxy)-1,2-benzenediamine

Using a procedure similar to Example 1B except starting with N¹-(*t*-butoxycarbonyl)-4-(dimethyl-*t*-butylsiloxy)-1,2-benzenediamine and 4-*t*-butylbenzoyl chloride, the title compound was obtained (97%); IR(CHCl₃): 858, 1271, 1514, 1668, 2962 cm⁻¹; NMR(300 MHz, CDCl₃): 0.22 (s, 6H); 0.98 (s, 9H); 1.35 (s, 9H); 1.55 (s, 9H); 6.46 (s, 1H); 6.64 (d, 1H, J=8.7 Hz); 7.06 (d, 1H, J=8.4 Hz); 7.47 (d, 2H, J=8.1 Hz); 7.88 (d, 2H, J=8.7 Hz); 8.93 (br s, 1H); MS(FD): 498.3.

- 59 -

Analysis for C₂₈H₄₂N₂O₄Si:

Calculated: C, 67.43; H, 8.49; N, 5.62

Found: C, 68.65; H, 8.82; N, 5.61

- 5 I) N¹-(t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-hydroxy-1,2-benzenediamine

Using a procedure similar to Example 1E except starting with N¹-(t-butoxy-carbonyl)-N²-(t-butylbenzoyl)-4-(dimethyl-t-butylsiloxy)-1,2-benzenediamine, the title compound was
10 obtained (97%); IR(KBr): 1168, 1285, 1536, 1631, 1690, 2965 cm⁻¹; NMR(300 MHz, CDCl₃): 1.36 (s, 9H); 1.53 (s, 9H); 6.38 (s, 1H); 6.66 (d, 1H, J=8.4 Hz); 7.01 (d, 1H, J=8.4 Hz); 7.49 (d, 2H, J=8.4 Hz); 7.70 (s, 1H); 7.89 (d, 2H, J=8.4 Hz); MS(FD): 384.2.

- 15 Analysis for C₂₂H₂₈N₂O₄ · 0.25 H₂O:

Calculated: C, 67.93; H, 7.39; N, 7.20

Found: C, 67.90; H, 7.37; N, 6.85

- 20 J) N¹-(4-t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine

Using a procedure similar to Example 2A except starting with N¹-(4-t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-hydroxy-1,2-benzenediamine and ethyl bromoacetate, the title
25 compound was obtained (76%); IR(CHCl₃): 1159, 1529, 1705, 1755, 2968 cm⁻¹; NMR(300 MHz, CDCl₃): 1.31 (t, 3H, J=6.9 Hz); 1.36 (s, 9H); 1.52 (s, 9H); 4.28 (q, 2H, J=6.9 Hz); 4.63 (s, 2H); 6.56 (s, 1H); 6.77 (d, 1H, J=8.7 Hz); 7.13 (d, 1H, J=8.7 Hz); 7.49 (d, 2H, J=8.4 Hz); 7.56 (s, 1H); 7.89 (d, 2H, J=8.1 Hz); 9.09 (br s, 1H); MS(FD): 470.

- 30 Analysis for C₂₆H₃₄N₂O₆:

Calculated: C, 66.36; H, 7.28; N, 5.95

Found: C, 68.08; H, 7.84; N, 5.54

- 60 -

K) N^2 -(*t*-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine

To a mixture of N^1 -(*t*-butoxycarbonyl)- N^2 -(*t*-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine (878 mg, 1.87 mmol) and CH_2Cl_2 (20 mL) was added TFA (1.5 mL, 19.5 mmol). After 12 h, the reaction was diluted with CH_2Cl_2 (300 mL) and washed with sat. aq. $NaHCO_3$ (3 x 50 mL). The organic layer was $MgSO_4$ dried, and concentrated. The crude material was chromatographed (65 g silica, 20% EtOAc/hexanes to 50% EtOAc/hexanes) to give the title compound (98%); IR($CHCl_3$): 1176, 1512, 1668, 1755, 2968 cm^{-1} ; NMR(300 MHz, $DMSO-d_6$): 1.22 (t, 3H, $J=7.2$ Hz); 1.33 (s, 9H); 4.17 (q, 2H, $J=7.2$ Hz); 4.62 (s, 2H); 6.64 (d, 1H, $J=8.7$ Hz); 6.74 (d, 1H, $J=8.7$ Hz); 6.95 (d, 1H, $J=3.0$ Hz); 7.54 (d, 2H, $J=8.7$ Hz); 7.91 (d, 2H, $J=8.1$ Hz); 9.59 (s, 1H); MS(FD): 370.

Analysis for $C_{21}H_{36}N_2O_4$:

Calculated: C, 68.09; H, 7.07; N, 7.56

Found: C, 68.34; H, 7.25; N, 7.56

L) N^1 -[4-(*N,N*-(bis-*t*-butoxycarbonyl)aminomethyl)-3-cyanobenzoyl]- N^2 -(*t*-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine

To a mixture of 4-(*N,N*-(bis-(*t*-butoxycarbonyl)aminomethyl)-3-cyanobenzoic acid (184 mg, 0.49 mmol) and CH_2Cl_2 (5 mL) was added, in sequence, pyridine (44 mL, 0.54 mmol), DMF (3 drops) and oxalyl chloride (48 mL, 0.55 mmol). After 1 h, the reaction was concentrated and the residue was diluted with CH_2Cl_2 (2 mL). Pyridine was added, followed by the N^2 -(*t*-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine (167 mg, 0.45 mmol) in CH_2Cl_2 (3 mL). After 1 h, the reaction was diluted with CH_2Cl_2 (75 mL) and washed with sat. aq. NH_4Cl (2 x 5 mL). The organic layer was $MgSO_4$ dried, filtered, and concentrated. The residue

- 61 -

chromatographed (30% EtOAc/hexanes) to give the title compound (140 mg, 0.19 mmol, 39%);

NMR(300 MHz, CDCl₃): 1.27 (t, 3H, J=6.6 Hz); 1.37 (s, 9H); 1.46 (s, 18H); 4.25 (q, 2H, J=7.2 Hz); 4.45 (s, 2H); 5.09 (s, 2H); 6.69 (d, 1H, J=9.0 Hz); 6.97 (s, 1H); 7.46 (m, 2H); 7.54 (d, 2H, J=8.7 Hz); 7.91 (d, 2H, J=8.4 Hz); 8.15 (d, 1H, J=8.7 Hz); 8.30 (s, 1H); 8.89 (s, 1H); 9.48 (s, 1H); MS(FD): 728.6.

Analysis for C₄₀H₄₈N₄O₉:

10 Calculated: C, 65.92; H, 6.64; N, 7.69
 Found: C, 66.33; H, 7.03; N, 7.39

M) N¹-[(1-iminoisoindolin-6-yl)carbonyl]-N²-(t-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine trifluoroacetate

15 To a mixture of N¹-[4-(N,N-(bis-t-butoxycarbonyl)-aminomethyl)-3-cyanobenzoyl]-N²-(t-butylbenzoyl)-4-(ethoxycarbonylmethoxy)-1,2-benzenediamine (109 mg, 0.15 mmol) in CH₂Cl₂ (5ml) was added TFA (0.15 mL, 1.95 mmol). After 1.5 h, more TFA (60 mL, 0.78 mmol) followed by TFA (1 mL, 12.98 mmol) were added. After 15 minutes, the reaction was concentrated and the residue was dissolved in EtOH (5 mL). Et₃N (42 mL, 0.30 mmol) was added. The solution was heated to reflux for 30 min. The reaction was cooled and concentrated. The residue was chromatographed with a polystyrene support (HP-20) column using H₂O as the eluant to give the title compound (29 mg, 0.045 mmol, 30%); NMR(300 MHz, DMSO-d₆): 1.30 (m, 12H); 4.29 (q, 2H, J=7.2 Hz); 4.65 (s, 2H); 4.67 (s, 2H); 6.97 (m, 2H); 7.45 (d, 2H, J=8.4 Hz); 7.52 (d, 1H, J=8.7 Hz); 7.65 (d, 1H, J=8.1 Hz); 8.00 (d, 2H, J=8.1 Hz); 8.23 (d, 1H, J=7.8 Hz); 8.49 (s, 1H); 9.48 (s, 2H); 9.62 (s, 1H); 10.07 (s, 1H); 10.20 (s, 1H); MS(FD): 529.3.

- 62 -

Example 9

Preparation of N¹-[(1-iminoisoindolin-6-yl)carbonyl]-N²-
(t-butylbenzoyl)-4-(benzyloxy)-1,2-benzenediamine
trifluoroacetate

5

A) N¹-(4-t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-benzyl-
oxy-1,2-benzenediamine

Using a procedure similar to Example 2A except starting
with N¹-(4-t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-hydroxy-
10 1,2-benzenediamine and benzyl bromide, the title compound
was obtained (65%); IR(CHCl₃): 1159, 1529, 1668, 1700 cm⁻¹;
NMR(300 MHz, DMSO-d₆): 1.33 (s, 9H); 1.44 (s, 9H); 5.10 (s,
2H); 6.87 (d, 1H, J=8.7 Hz); 7.41 (m, 7H); 7.57 (d, 2H, J-
8.4 Hz); 7.87 (d, 2H, J=8.4 Hz); 8.51 (br s, 1H); 9.77 (s,
15 1H); MS(FD): 474.

Analysis for C₂₉H₃₄N₂O₄:

Calculated: C, 73.39; H, 7.22; N, 5.90

Found: C, 74.29; H, 7.92; N, 5.48

20 B) N²-(t-butylbenzoyl)-4-benzyloxy-1,2-benzenediamine

Using a procedure similar to Example 8K except starting
with N¹-(4-t-butoxycarbonyl)-N²-(t-butylbenzoyl)-4-benzyl-
oxy-1,2-benzenediamine, the title compound was obtained
(72%); IR(CHCl₃): 1270, 1511, 1609, 1667, 2967 cm⁻¹;

25 NMR(300 MHz, DMSO-d₆): 1.33 (s, 9H); 4.99 (s, 2H); 6.74 (m,
3H); 7.03 (s, 1H); 7.39 (m, 6H); 7.54 (d, 2H, J=8.4 Hz);
7.92 (d, 2H, J=8.7 Hz); 9.59 (s, 1H); MS(FD): 374.1

Analysis for C₂₄H₂₆N₂O₂:

Calculated: C, 76.98; H, 7.00; N, 7.48

30 Found: C, 77.02; H, 6.76; N, 7.27

C) N¹-[4-(N,N-(bis-t-butoxycarbonyl)aminomethyl)-3-cyano-
benzoyl]-N²-(t-butylbenzoyl)-4-(benzyloxy)-1,2-benzene-
diamine

- 63 -

Using a procedure similar to Example 8L except starting with N²-(t-butylbenzoyl)-4-benzyloxy-1,2-benzenediamine, the title compound was obtained (81%).

NMR(300 MHz, CDCl₃): 1.33 (s, 9H); 1.46 (s, 18H); 4.89 (s, 2H); 5.07 (s, 2H); 6.75 (d, 1H, J=9.0 Hz); 7.03 (s, 1H); 7.40 (m, 6H); 7.55 (d, 2H, J=8.4 Hz); 7.92 (d, 2H, J=8.4 Hz); 8.14 (d, 1H, J=8.4 Hz); 8.28 (s, 1H); 8.82 (s, 1H); 9.39 (s, 1H); MS(FD): 732.3.

Analysis for C₄₃H₄₈N₄O₇:

Calculated: C, 70.47; H, 6.60; N, 7.65
Found: C, 70.52; H, 6.64; N, 7.70

D) N¹-[(1-iminoisoindolin-6-yl)carbonyl]-N²-(4-t-butylbenzoyl)-4-benzyloxy-1,2-benzenediamine trifluoroacetate

To a mixture of N¹-[4-(N,N-(bis-t-butoxycarbonyl)-aminomethyl)-3-cyanobenzoyl]-N²-(t-butylbenzoyl)-4-benzyloxy-1,2-benzenediamine (176 mg, 0.24 mmol) and CH₂Cl₂ (5 mL) was added TFA (1 mL, 12.98 mmol). After 1 h, the reaction was concentrated and the residue was dissolved in EtOH (5 mL) and heated to reflux. After 1 h, the reaction was cooled and concentrated to give the title compound as a tan solid (156 mg, 0.24 mmol, 100%); IR(KBr): 1202, 1462, 1500, 1664, 1696cm⁻¹; NMR(300 MHz, DMSO-d₆): 1.26 (s, 9H); 4.85 (s, 2H); 5.12 (s, 2H); 6.96 (d, 1H, J=8.7 Hz); 7.42 (m, 9H); 7.82 (d, 2H, J=8.4 Hz); 7.87 (d, 1H, J=8.1 Hz); 8.30 (d, 1H, J=8.1 Hz); 9.37 (s, 2H); 9.75 (s, 1H); 9.89 (s, 1H); 10.13 (s, 1H); MS(FD): 533.3.

Analysis for C₃₃H₃₂N₄O₃·CF₃COOH·0.25 H₂O:

Calculated: C, 64.56; H, 5.19; N, 8.60
Found: C, 64.28; H, 5.58; N, 7.60

- 64 -

Example 10

Preparation of N¹-[(1-iminoisoindolin-6-yl)carbonyl]-N²-(t-butylbenzoyl)-4-(hydroxy)-1,2-benzenediamine trifluoroacetate

5

A) N¹-[4-(N,N-(bis-t-butoxycarbonyl)aminomethyl)-3-cyano-benzoyl]-N²-(t-butylbenzoyl)-4-(hydroxy)-1,2-benzenediamine

Using a procedure similar to Example 1C except starting

10 with N¹-[4-(N,N-(bis-t-butoxycarbonyl)aminomethyl)-3-cyano-benzoyl]-N²-(t-butylbenzoyl)-4-(benzyloxy)-1,2-benzene-diamine, the title compound was obtained (92%); IR(CHCl₃): 1233, 1370, 1649, 1749, 2970 cm⁻¹; NMR(300 MHz, CDCl₃): 5.08 (s, 2H); 6.39 (d, 1H, J=6.9 Hz); 7.16 (d, 1H, J=6.9 Hz); 7.21 (s, 1H); 7.35 (d, 1H, J=8.7 Hz); 7.54 (d, 2H, J=8.1 Hz); 7.88 (d, 2H, J=8.1 Hz); 8.22 (d, 1H, J=8.7 Hz); 8.37 (s, 1H); 9.44 (s, 1H); 9.52 (s, 1H); MS(FD): 542.1 (loss of BOC).

Analysis for C₃₆H₄₂N₄O₇:

Calculated: C, 67.27; H, 6.59; N, 8.72

20 Found: C, 67.50; H, 6.67; N, 8.55

B) N¹-[(1-iminoisoindolin-6-yl)carbonyl]-N²-(t-butylbenzoyl)-4-hydroxy-1,2-benzenediamine trifluoroacetate

Using a procedure similar to Example 9 except starting

25 with N¹-[4-(N,N-(bis-t-butoxycarbonyl)aminomethyl)-3-cyano-benzoyl]-N²-(t-butylbenzoyl)-4-hydroxy-1,2-benzenediamine, the title compound was obtained as a tan solid (quant.); IR(KBr): 1135, 1202, 1512, 1670 cm⁻¹; NMR(300 MHz, DMSO-d₆): 1.24 (s, 9H); 4.84 (s, 2H); 6.66 (d, 1H, J=8.7 Hz); 7.17 (s, 1H); 7.29 (d, 1H, J=8.7 Hz); 7.46 (d, 2H, J=8.1 Hz); 7.79 (d, 2H, J=8.1 Hz); 7.85 (d, 1H, J=8.1 Hz); 8.27 (d, 1H, J=7.8 Hz); 8.79 (s, 1H); 9.33 (s, 1H); 9.78 (s, 1H); 10.05 (s, 1H); 10.51 (s, 1H); MS(FD): 443.2.

30 Analysis for C₂₆H₂₆N₄O₃·CF₃COOH·2H₂O

- 65 -

Calculated: C, 56.76; H, 5.27; N, 9.45

Found: C, 56.71; H, 4.91; N, 8.67

Example 11**5 Preparation of 2-(4-t-butylbenzoylamino)-N-[3-(amino-(imino)methyl)phenyl]benzamide hydrochloride****A) 2-nitro-N-(3-cyanophenyl)benzamide**

Using a procedure similar to Example 1D except starting
10 with 2-nitrobenzoic and 3-cyanoaniline, the title compound
was obtained as a solid (92%); IR(KBr): 1344, 1524, 1645,
1666, 2229, 3232 cm^{-1} ; NMR(300 MHz, CDCl_3): 7.49 (m, 2H);
7.71 (m, 3H); 7.81 (m, 2H); 7.99 (s, 1H); 8.18 d, 1H, J=8.1
Hz); MS(FD): 267.1.

15 Analysis for $\text{C}_{14}\text{H}_9\text{N}_3\text{O}_3$:

Calculated: C, 62.92; H, 3.40; N, 15.72

Found: C, 62.74; H, 3.52; N, 15.71

B) 2-amino-N-(3-cyanophenyl)benzamide

20 Using a procedure similar to Example 1C except starting
with 2-nitro-N-(3-cyanophenyl)benzamide, the title compound
was obtained as a solid (97%); IR(CHCl_3): 1527, 1583, 1614,
1668, 2234 cm^{-1} ; NMR(300 MHz, CDCl_3): 6.76 (d, 1H, J=6.9
Hz); 7.32 (d, 1H, J=7.5 Hz); 7.48 (m, 4H); 7.76 (d, 1H,
25 J=7.8 Hz); 7.88 (br s, 1H); 8.05 (s, 1H); MS(FD): 237.1.

Analysis for $\text{C}_{14}\text{H}_{11}\text{N}_3\text{O} \cdot 0.33\text{H}_2\text{O}$:

Calculated: C, 69.30; H, 4.82; N, 17.32

Found: C, 69.61; H, 4.87; N, 17.16

30 C) 2-(4-t-butylbenzoylamino)-N-(3-cyanophenyl)benzamide

Using a procedure similar to Example 1B except starting
with 2-amino-N-(3-cyanophenyl)benzamide and 4-t-butylbenzoyl
chloride, the title compound was obtained as a white solid
(92%); MP: 194-6 °C; IR(KBr): 757, 1584, 1647, 2969, 3256

- 66 -

cm⁻¹; NMR(300 MHz, CDCl₃): 1.40 (s, 9H); 6.90 (t, 1H, J=7.2 Hz); 7.31 (m, 1H); 7.49-7.55 (m, 3H); 7.60 (d, 2H, J=7.5 Hz); 7.95 (d, 3H, J=8.4 Hz); 8.36 (s, 1H); 8.46 (d, 1H, J=8.4 Hz); 9.19 (s, 1H); 11.42 (s, 1H); MS(FD): 397.1.

5 Analysis for C₂₅H₂₃N₃O₂·0.25H₂O

Calculated: C, 74.70; H, 5.89; N, 10.45

Found: C, 74.84; H, 6.07; N, 9.60

10 D) 2-(4-t-butylbenzoylamino)-N-[3-(amino(hydroxyimino)-methyl)phenyl]benzamide

Using a procedure similar to Example 1F except starting with 2-(4-t-butylbenzoylamino)-N-(3-cyanophenyl)benzamide, the title compound was obtained as a white solid (42%);

15 NMR(300 MHz, CDCl₃): 1.34 (s, 9H); 5.02 (br s, 2H); 7.08 (t, 1H, J=2.1 Hz); 7.35-7.45 (m, 2H); 7.49 (d, 2H, J=6.9 Hz); 7.69 (d, 1H, J=6.0 Hz); 7.74 (d, 1H, J=8.3 Hz); 7.84 (s, 1H); 7.92 (d, 2H, J=8.1 Hz); 8.70 (d, 2H, J=8.4 Hz); 8.83 (s, 1H); 11.66 (s, 1H); MS(FD): 430.2.

Analysis for C₂₅H₂₆N₄O₃·H₂O:

20 Calculated: C, 66.95; H, 6.29; N, 12.49

Found: C, 67.34; H, 5.88; N, 12.31

E) 2-(4-t-butylbenzoylamino)-N-[3-(amino(imino)methyl)-phenyl]benzamide hydrochloride

25 Using a procedure similar to Example 1G except starting with 2-(4-t-butylbenzoyl-amino)-N-3-(amino(hydroxyimino)-methyl)phenyl]benzamide, the title compound was obtained as a tan solid (66%); IR(CHCl₃): 1528, 1656, 1676, 2968 cm⁻¹; NMR(300 MHz, DMSO-d₆): 1.29 (s, 9H); 7.30 (t, 1H, J=7.2 Hz); 7.57 (m, 5H); 7.83 (d, 2H, J=7.8 Hz); 7.89 (d, 1H, J=7.5 Hz); 7.94 (d, 1H, J=8.1); 8.17 (s, 1H); 8.36 (d, 1H, J=8.1 Hz); 9.01 (s, 2H); 9.35 (s, 2H); 10.85 (s, 1H); 11.42 (s, 1H); MS(FD): 415.3.

Analysis for C₂₅H₂₇N₄O₂·HCl·1.5H₂O

- 67 -

Calculated: C, 62.82; H, 6.33; N, 11.72

Found: C, 62.36; H, 6.36; N, 11.60

Example 12**5 Preparation of 2-[3-(amino(imino)methyl)benzoylamino]-N-(4-methoxyphenyl)benzamide****A) 2-amino-N-(4-methoxyphenyl)benzamide**

A mixture of isatoic anhydride (4.894 g, 30 mmol),
10 4-methoxyaniline (3.695 g, 30 mmol) and toluene (60 mL) was
heated to reflux for 5 h. After cooling, the reaction was
filtered. The solid was triturated with CH₂Cl₂ (500 mL).
This supernatant was combined with the above filtrate and
concentrate. The residue was dissolved in CH₂Cl₂,
15 decolorized with activated charcoal and crystallized from
CH₂Cl₂/hexanes to give the title compound as a white solid
(5.3 g, 73%); mp 116-7 C; MS(FD): 242.

Analysis for C₁₄H₁₄N₂O₂:

Calculated: C, 69.41; H, 5.83; N, 11.56

20 Found: C, 69.16; H, 5.91; N, 11.31

B) 2-(3-cyanobenzoylamino)-N-(4-methoxyphenyl)benzamide

Using a procedure similar to Example 1B except starting
with 2-amino-N-(4-methoxyphenyl)benzamide, the title
25 compound was obtained as a white solid (77%); mp 190-2 C;
3.74 (s, 3H), 6.94 (d, 2H, J=12.3 Hz), 7.33 (dd, 1H, J=8.2,
8.2 Hz), 7.60 (2H, J=12.3 Hz), 7.62 (m, 1H), 7.80 (dd, 1H,
J=10.2, 8.2 Hz), 7.90 (d, 1H, J=8.2 Hz), 8.10 (d, 1H, J=8.2
Hz), 8.20 (d, 1H, J=8.2 Hz), 8.30 (s, 1H), 8.35 (1H, J=8.2
30 Hz), 10.43 (s, 1H), 11.75 ((s, 1H); MS(FD): 371.

Analysis for C₂₂H₁₇N₃O₃·H₂O:

Calculated: C, 67.86; H, 4.92; N, 10.79

Found: C, 68.10; H, 4.88; N, 10.85

- 68 -

C) 2-[3-(amino(hydroxyimino)methyl)benzoylamino]-N-(4-methoxyphenyl)benzamide

Using a procedure similar to Example 1F except starting with 2-(3-cyanobenzoylamino)-N-(4-methoxyphenyl)benzamide, the title compound was obtained as a white solid (98%); mp 204-7 C; MS(FD): 404 (M+H).

Analysis for C₂₂H₂₀N₄O₄:

Calculated: C, 65.34; H, 4.99; N, 13.85

Found: C, 66.00; H, 4.96; N, 13.54

D) 2-[3-(amino(imino)methyl)benzoylamino]-N-(4-methoxyphenyl)benzamide

Using a procedure similar to Example 1G except starting with 2-[3-(amino(hydroxyimino)methyl)benzoylamino]-N-(4-methoxyphenyl)benzamide, the title compound was obtained as a light yellow solid (54%); mp 163-7 C; MS(FD): 388.1.

Analysis for C₂₂H₂₁N₄O₃·HCl·2.5 H₂O:

Calculated: C, 56.23; H, 5.58; N, 11.92

Found: C, 55.91; H, 5.07; N, 11.71

Example 13

Preparation of 3-(4-tert-butylbenzoylamino)-N-[3-(amino(imino)-methyl)phenyl]thiophene-2-carboxamide hydrochloride

A) methyl 3-(4-tert-butylbenzoylamino)thiophene-2-carboxylate

To a mixture of methyl 3-aminothiophene-2-carboxylic acid (5.0g, 35.0 mmol), pyridine (2.82 mL, 35.0 mmol), and dry methylene chloride (160 mL) at 0 C was added 4-tert-butylbenzoyl chloride (6.21 mL, 31.8 mmol) and stirred for 1 h. The reaction was concentrated, dissolved in ethyl acetate, washed with water (4X) and brine, MgSO₄ dried and concentrated. The residue was chromatographed (2% ethyl acetate/hexanes to 5% ethyl acetate/hexanes) to afford the title compound as a solid (9.67 g, 96% yield); NMR(CDCl₃) δ

- 69 -

1.36(s, 9H), 3.93(s, 3H), 7.54(d, J=8.4 Hz, 2H), 7.54(d, J=5.4 Hz, 1H), 7.95(d, J=8.4 Hz, 2H), 8.30(d, J=5.4 Hz, 1H), 11.16(s, 1H); MS(FD): 317.

Analysis for C₁₇H₁₉NO₈S:

5 Calculated: C, 64.33; H, 6.03; N, 4.42
 Found: C, 64.39; H, 5.98; N, 4.46.

B) 3-(4-t-butylbenzoylamino)thiophene-2-carboxylic acid

10 A mixture of methyl 3-(4-t-butylbenzoylamino)thiophene-2-carboxylate (9.67 g, 30 mmol), dioxane (75 mL) and 2 M sodium hydroxide (75 mL) was stirred for 16 h. The reaction was acidified to pH 2 with 5 M hydrochloric acid. The mixture was diluted with ethyl acetate and the aqueous layer
15 was extracted with ethyl acetate (3 times). The combined organic layers was magnesium sulfate dried, filtered, and concentrated to afford the title compound as a solid (8.09 g, 89%); NMR(CDCl₃) d 1.36(s, 9H), 7.54(d, 2H, J=8.4 Hz), 7.61(d, 1H, J=5.1 Hz), 7.92(d, 2H, J=8.4 Hz), 8.34(d, 1H, J=5.1 Hz), 11.04(s, 1H); MS(FD): 303.

Analysis for C₁₆H₁₇NO₈S:

 Calculated: C, 63.34; H, 5.93; N, 4.62.
 Found: C, 63.56; H, 5.93; N, 4.32.

25

C) 2-(4-t-butylphenyl)-4H-thieno[3,2-d][1,3]oxazin-4-one

 To a mixture of 3-(4-t-butylbenzoylamino)thiophene-2-carboxylic acid (8.09 g, 27 mmol) and methylene chloride (135 mL) was added oxalyl chloride (11.8 mL, 135 mmol). The
30 mixture was heated until bubbling began. The reaction was stirred for 2 h without heating. The reaction was concentrated and the residue was dissolved in methylene chloride (135 mL). To this solution was added pyridine (2.2 mL, 27 mmol) and stirred at room temperature for 1 h.

- 70 -

The reaction mixture was concentrated and partitioned between ethyl acetate and water. The organic layer was washed with water (4 times) and brine, MgSO_4 dried and concentrated. The residue was chromatographed (10% ethyl acetate/hexanes) to afford the title compound as a solid (7.44 g, 96%); NMR(CDCl_3) δ 1.37(s, 9H), 7.35(d, J=5.1 Hz, 1H), 7.52(d, J=8.4 Hz, 2H), 7.91(d, J=5.1 Hz, 1H), 8.22(d, J=8.4 Hz, 2H); MS(FD): 285.

Analysis for $\text{C}_{16}\text{H}_{15}\text{NO}_2\text{S}$:

10 Calculated: C, 67.34; H, 5.30; N, 4.91.
 Found: C, 67.51; H, 5.56; N, 4.76.

D) 3-(4-t-butylbenzoylamino)-N-(3-cyanophenyl)thiophene-2-carboxamide

15 To a mixture of 2-(4-t-butylphenyl)-4H-thieno[3,2-d]-[1,3]oxazin-4-one (400 mg, 1.4 mmol), 3-aminobenzonitrile (165 mg, 1.4 mmol) and THF (5 mL) was added 0.5 M potassium hexamethyldisilazide in toluene (3.08 mL, 1.54 mmol) and stirred for 15 min. The reaction was added NH_4Cl and ethyl
20 acetate. The organic layer was washed with water (3X) and brine, MgSO_4 dried and concentrated. The residue was chromatographed (20% THF/hexanes) to give the title compound as a solid (580 mg, 51%); 1.37 (s, 9H), 7.47 (m, 4H), 7.57 (d, 2H, J=8.4 Hz); 7.68 (m, 1H), 7.98 (d, 2H, J=8.4 Hz);
25 8.20 (s, 1H); 8.42 (d, 1H, J=5.7 Hz); 11.80 (s, 1H); MS(FD): 403.

Analysis for $\text{C}_{23}\text{H}_{21}\text{N}_3\text{O}_2\text{S}$:

 Calculated: C, 68.46; H, 5.25; N, 10.41
 Found: C, 67.83; H, 5.14; N, 10.08.

30

E) 3-(4-t-butylbenzoylamino)-N-[3-(amino(imino)methyl)-phenyl]thiophene-2-carboxamide hydrochloride

- 71 -

Into a mixture of 3-(4-t-butylbenzoylamino)-N-(3-cyano-phenyl)thiophene-2-carboxamide (400 mg, 0.993 mmol), pyridine (10 mL), and triethylamine (1 mL) was bubbled hydrogen sulfide gas for 10 min. The solution was allowed to stand overnight. The solvent was concentrated and the residue was dissolved in ethyl acetate and washed with water (2X) and saturated sodium bicarbonate. The organic layer was MgSO₄ dried, filtered, and concentrated. The residue was dissolved in a mixture of 1:1 iodomethane : methanol (10 mL) and refluxed for 30 min. The reaction was concentrated and the residue was vacuum dried for 1 h. This residue was dissolved in a minimum amount of dry methanol and treated with ammonium acetate (230 mg, 2.98 mmol). The solution was heated to 65 degrees for 1 h, cooled to room temperature. The solution was triturated with diethyl ether and the solid was filtered off. The solid was dissolved in 10 mL of 1:1 tetrahydrofuran/water and treated with di-tert-butyl dicarbonate (1.08g, 4.96 mmol) and potassium carbonate (822 mg, 5.96 mmol). After being stirred vigorously for 30 minutes, the solution was partitioned between with ethyl acetate and water. The aqueous layer was extracted with ethyl acetate (2X). The combined organic phase was MgSO₄ dried and concentrated. The residue was chromatographed (30% EtOAc/CH₂Cl₂) to give 3-(4-t-butylbenzoylamino)-N-[3-(N-t-butoxycarbonylamino(N-t-butoxycarbonylimino)methyl)phenyl]thiophene-2-carboxamide. This material was dissolved in trifluoroacetic acid (2 mL) and stirred for 1 h. The reaction was concentrated and the residue was dissolved in ethyl acetate. To this solution was added 1 M hydrochloric acid in ether, and the resulting mixture was sonicated for 5 min. The resulting solid was filtered, dissolved in water, and lyophilized to afford the title compound as a powder (170mg, 37% yield); NMR(DMSO-d)
δ 1.38(s, 9H), 7.56 (m, 1H), 7.63 (m, 3H), 7.83 (d, 2H,

- 72 -

J=8.29 Hz), 7.96 (d, 1H, J=5.27 Hz), 8.05 (m, 1H), 8.16 (m, 2H); MS(FD): 421.

Analysis for $C_{23}H_{24}N_4O_2S \cdot HCl \cdot 0.8 H_2O$:


Calculated: C, 58.61; H, 5.69; N, 11.89

5 Found: C, 58.31; H, 5.40; N, 11.84.

Example 14

Preparation of N^1 -[3-[amino(hydroxyimino)methyl]benzoyl]]-
 N^2 -(4-methoxybenzoyl)-1,2-benzenediamine

10



A) N-(4-methoxybenzoyl)-2-nitroaniline

Using a procedure similar to Example 1B except starting
15 with 2-nitroaniline and p-anisoyl chloride, the title
compound was obtained as a solid (98%); MS(FD): 272.0.

Analysis for $C_{14}H_{12}N_2O_4$:

Calculated: C, 61.76; H, 4.44; N, 10.29

Found: C, 61.87; H, 4.61; N, 9.99

20

B) N^1 -(4-methoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1C except starting
with N-(4-methoxybenzoyl)-2-nitroaniline in ethyl acetate,
the title compound was obtained as a white solid (99%);
25 MS(FD): 242.

Analysis for $C_{14}H_{14}N_2O_2$:

Calculated: C, 69.41; H, 5.82; N, 11.56

Found C, 69.63; H, 5.87; N, 11.39

- 73 -

C) N¹-(3-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1B except starting with N¹-(4-methoxybenzoyl)-1,2-benzenediamine, the title
5 compound was obtained as a solid (95%).

Analysis for C₂₂H₁₇N₃O₃•0.5H₂O: Calculated: C, 69.46; H, 4.77; N, 11.05

Found: C, 69.53; H, 4.57; N, 10.82

10 D) N¹-[3-[amino(hydroxyimino)methyl]benzoyl]]-N²-(4-methoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting with N¹-(3-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid
15 (18%); MS(FD): 404.

Analysis for C₂₂H₂₀N₄O₄:


Calculated: C, 65.34; H, 4.98; N, 13.85;

Found: C, 65.29; H, 5.06; N, 13.83

20

Example 15

Preparation of N¹-[3-(aminoiminomethyl)benzoyl]]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride



25

A solution of N¹-(3-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine (0.5 g, 1.35 mmol) in triethylamine (5 mL) and pyridine (50 mL) was bubbled with hydrogen sulfide for 5 min. The reaction was stoppered and allowed
30 to stand for 48 h. The reaction was concentrated and the

- 74 -

residue was dissolved in ethyl acetate and washed twice with brine. The organic phase was MgSO_4 dried, filtered, and concentrated. The residue was dissolved in acetone (100 mL) and iodomethane (25 mL, 641.7 mmol) was added. The reaction
5 was heated to reflux for 2 h, cooled, and concentrated. The residue was then dissolved in methanol (100 mL) and ammonium acetate (0.15 g, 5.4 mmol) was added. This solution was heated to reflux for 12 h, cooled and concentrated. The resulting residue was dissolved in THF (20 mL) and di-
10 t-butyl dicarbonate (0.87 g, 4 mmol) was added, followed by a solution of K_2CO_3 (0.93 g, 6.75 mmol) in water (10 mL). After 2 h, the reaction was concentrated. The residue was dissolved in a minimal amount of chloroform and chromatographed (30% ethyl acetate/hexanes to ethyl acetate)
15 to give the title compound as the free base. This material was dissolved in trifluoroacetic acid, stirred for 2 h, and concentrated. To the residue was added 1N HCl, and the mixture was stirred vigorously, filtered and vacuum dried to give the title compound as a white solid (220 mg, 42%);
20 MS(FD): 389.1 (MH^+).

Analysis for $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_3 \cdot \text{HCl}$:


Calculated: C, 62.19; H, 4.98; N, 13.19; Cl, 8.34

Found C, 61.92; H, 5.09; N, 13.30; Cl, 8.22

25

Example 16

Preparation of N^1 -[4-[amino(hydroxyimino)methyl]benzoyl]- N^2 -(4-methoxybenzoyl)]-1,2-benzenediamine



30

- 75 -

A) N¹-(4-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1B except starting with N¹-(4-methoxybenzoyl)-1,2-benzenediamine and 4-cyanobenzoyl chloride, the title compound was obtained as a solid (59%); MS(FD): 371.0.

Analysis for C₂₂H₁₇N₃O₃:

Calculated: C, 71.15; H, 4.61; N, 11.31

Found C, 70.94; H, 4.85; N, 11.05

B) N¹-[4-[amino(hydroxyimino)methyl]benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting with N¹-(4-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (19%); MS(FD): 404.1.


Analysis for C₂₂H₂₀N₄O₄:

Calculated: C, 65.34; H, 4.99; N, 13.85

Found C, 64.87; H, 4.97; N, 13.31

Example 17

Preparation of N¹-[4-(aminoiminomethyl)benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride



Using a procedure similar to Example 1G except starting with N¹-[4-[amino(hydroxyimino)methyl]benzoyl]-N²-

- 76 -

(4-methoxybenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (26%); MS(FD): 389.1 (MH⁺).

Analysis for C₂₂H₂₀N₄O₃·1.1HCl·0.5H₂O:


Calculated: C, 60.39; H, 5.09; N, 12.80

5 Found: C, 60.30; H, 4.90; N, 12.48

Example 18

Preparation of N¹-[2-fluoro-4-(aminoiminomethyl)benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride

10



A) N¹-(2-fluoro-4-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine

15 To a solution of N¹-(4-methoxybenzoyl)-1,2-benzenediamine (0.5 g, 1.8 mmol), 2-fluoro-4-cyanobenzoic acid (0.3 g, 1.8 mmol) and N,N-diisopropylethylamine (0.23 g, 1.8 mmol) in DMF (5 mL) was added EDC (0.42 g, 2.2 mmol). After 48 h, 2-fluoro-4-cyanobenzoic acid (0.3 g, 1.8 mmol) and EDC
20 (0.35 g, 1.8 mmol) were added. After an additional 24 h, the reaction was concentrated; and the residue was dissolved in ethyl acetate, washed with 1 N citric acid, brine, satd aq NaHCO₃ and brine. The organic layer was MgSO₄ dried, crystallized, filtered with ether wash, and dried under
25 vacuum to give the title compound as a white solid 280 mg (40%); MS(FD): 389.1

Analysis for C₂₂H₁₆FN₃O₃:

Calculated: C, 67.86; H, 4.14; N, 10.79

Found C, 68.07; H, 4.20; N, 10.66

- 77 -

B) N^1 -[2-fluoro-4-(aminoiminomethyl)benzoyl]- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 15 except starting
5 with N^1 -(2-fluoro-4-cyanobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine, the title compound as obtained as a solid (12%); MS(ES): 407.1 (MH^+).


Analysis for $C_{22}H_{19}FN_4O_3 \cdot HCl$:

Calculated: C, 59.67; H, 4.55; N, 12.65

10 Found C, 59.45; H, 4.63; N, 12.39

Example 19

Preparation of 5-benzyloxycarbonyl- N^1 -[3-[amino-
(hydroxyimino)methyl]benzoyl]- N^2 -(4-*t*-butylbenzoyl)-1,2-
15 benzenediamine



20 A) 5-methoxycarbonyl- N^1 -(3-cyanobenzoyl)-1,2-benzenediamine

To a solution of 4-methoxycarbonyl-1,2-benzenediamine (8 g, 48 mmol) and pyridine (4 mL, 48 mmol) in acetonitrile (300 mL) at 0 °C was added 3-cyanobenzoyl chloride (8 g,
25 48 mmol). After 12 h, the reaction was concentrated and the residue was suspended with vigorous stirring in ethyl acetate (500 mL) and 1 M citric acid (200 mL). The solid was filtered, washed with diethyl ether and vacuum dried to

- 78 -

give the title compound as an off-white solid (8 g, 56%);
MS(FD): 295.1 (M^+).

Analysis for $C_{16}H_{13}N_3O_3$:

Calculated: C, 65.08; H, 4.44; N, 14.23

5 Found: C, 65.33; H, 4.54; N, 14.39

B) 5-methoxycarbonyl- N^1 -(3-cyanobenzoyl)- N^2 -(4-t-butylbenzoyl)]-1,2-benzenediamine

Using a procedure similar to Example 1B except starting
10 with 4-t-butylbenzoyl chloride and 5-methoxycarbonyl- N^1 -(3-cyanobenzoyl)]-1,2-benzenediamine, the title compound was obtained as a solid (2.32 g, 60%); MS(FD): 455.2.

Analysis for $C_{27}H_{25}N_3O_4$:

Calculated: C, 71.19; H, 5.53; N, 9.22

15 Found: C, 71.31; H, 5.82; N, 9.35

C) 5-benzyloxycarbonyl- N^1 -(3-cyanobenzoyl)- N^2 -(4-t-butylbenzoyl)]-1,2-benzenediamine

To a solution of 5-methoxycarbonyl- N^1 -(3-cyano-
20 benzoyl)]- N^2 -(4-t-butylbenzoyl)]-1,2-benzenediamine (2.6 g, 5.7 mmol) in THF (80 mL) was added methanol (27 mL), followed by a solution of $LiOH \cdot H_2O$ (1 g, 22.8 mmol) in water (27 mL). After 12 h, the pH was adjusted to 3 with conc HCl and the mixture was concentrated. The residue was
25 partitioned between ethyl acetate and water. The organic layer was washed with 1 N HCl, brine (2X), $MgSO_4$ dried, and concentrated. To the residue dissolved in DMF (20 mL) was added benzyl bromide (1.06 g, 6.2 mmol), followed by K_2CO_3 (1.6 g, 11.4 mmol). After 3 h, the reaction was
30 concentrated and the residue was dissolved in ethyl acetate and washed with satd aq $NaHCO_3$ followed by brine. The organic layer was $MgSO_4$ dried and 5 g of silica gel was added. This mixture was concentrated and chromatographed (20% ethyl acetate/hexanes to 60% ethyl acetate/hexanes) to

- 79 -

give the title compound as a white solid (1.2 g, 40%);
MS(FD): 531.0.

Analysis for $C_{33}H_{29}N_3O_4$:

Calculated: C, 74.56; H, 5.50; N, 7.90

5 Found: C, 74.32; H, 5.64; N, 7.63

D) 5-benzyloxycarbonyl- N^1 -[3-[amino(hydroxyimino)methyl]-benzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine

10 Using a procedure similar to Example 1F except starting with 5-benzyloxycarbonyl- N^1 -(3-cyanobenzoyl)- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (47%); MS(FD): 564 (M^+)

Analysis for $C_{33}H_{32}N_4O_5$:


Calculated: C, 69.10; H, 5.80; N, 9.76

15 Found: C, 69.52; H, 5.84; N, 9.42

Example 20

Preparation of 5-carboxy- N^1 -[3-(aminoiminomethyl)benzoyl]]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride

20



Using a procedure similar to Example 1G, except starting with 5-benzyloxycarbonyl- N^1 -[3-[amino-(hydroxyimino)methyl]benzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (330 mg, 83%).

Analysis for $C_{26}H_{26}N_4O_4 \cdot 1.1HCl \cdot 0.5H_2O$:


- 80 -

Calculated: C, 61.52; H, 5.58; N, 11.03; Cl, 7.68

Found: C, 61.15; H, 5.49; N, 11.09; Cl, 7.90.

Example 21

- 5 **Preparation of 4-benzyloxycarbonyl-N¹-[3-[amino-(hydroxyimino)methyl]benzoyl]]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine**



10

- A) 4-methoxycarbonyl-N²-(4-t-butylbenzoyl)-1,2-benzene-diamine

Using a procedure similar to Example 1B except starting with 4-methoxycarbonyl-1,2-benzenediamine and 4-t-butyl-benzoyl chloride, the title compound was obtained as a solid (10.8 g, 65%); MS(FD): 326.2.

Analysis for C₁₉H₂₂N₂O₃:

Calculated: C, 69.92; H, 6.79; N, 8.58

Found: C, 70.02; H, 6.91; N, 8.61

20

- B) 4-methoxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)]-1,2-benzenediamine

Using a procedure similar to Example 1B except starting with 3-cyanobenzoyl chloride and 4-methoxycarbonyl-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (3.4 g, 81%); MS(FD): 455.3.

Analysis for C₂₇H₂₅N₃O₄:

Calculated: C, 71.19; H, 5.53; N, 9.22

Found: C, 71.08; H, 5.59; N, 9.34

- 81 -

C) 4-benzyloxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 19 except starting
5 from 4-methoxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (3.3 g, 65%); MS(FD): 531.0.

Analysis for C₃₃H₂₉N₃O₄:

Calculated: C, 74.56; H, 5.50; N, 7.90

10 Found: C, 74.42; H, 5.73; N, 7.73

D) 4-benzyloxycarbonyl-N¹-[3-[amino(hydroxyimino)methyl]benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 1F except starting
15 from 4-benzyloxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (1.62 g, 76%); MS(FD): 564.

Analysis for C₃₃H₃₂N₄O₅:


Calculated: C, 70.70; H, 5.71; N, 9.92

20 Found: C, 70.25; H, 5.62; N, 9.81

Example 22

Preparation of 4-carboxy-N¹-[4-(aminoiminomethyl)benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride

25



Using a procedure similar to Example 1G except starting
from 4-benzyloxycarbonyl-N¹-[3-[amino(hydroxyimino)methyl]-

- 82 -

benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (394 mg, 90%); MS(FD): 459. Analysis for C₂₆H₂₆N₄O₄·1.15HCl·1H₂O:


Calculated: C, 60.24; H, 5.67; N, 10.80; Cl, 7.86

5 Found: C, 60.43; H, 5.46; N, 10.84; Cl, 7.84

Example 23

Preparation of 4-methoxycarbonyl-N¹-[3-[amino(hydroxyimino)-methyl]benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

10



Using a procedure similar to Example 1F except starting from 4-methoxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (1.6 g, 75%); MS(FD): 488.

15 Analysis for C₂₇H₂₈N₄O₅:

Calculated: C, 66.38; H, 5.78; N, 11.47

Found: C, 66.42; H, 5.91; N, 11.43


20

Example 24

Preparation of 4-methoxycarbonyl-N¹-[3-(aminoiminomethyl)-benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride

25


- 83 -



Using a procedure similar to Example 1G except starting from 4-methoxycarbonyl-N¹-[3-[amino(hydroxyimino)methyl]-benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (0.95 g, 57%); MS(FD): 473.

Example 25

10 **Preparation of 5-methoxycarbonyl-N¹-[3-[amino(hydroxyimino)methyl]benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine**



15 Using a procedure similar to Example 1F except starting from 5-methoxycarbonyl-N¹-(3-cyanobenzoyl)-N²-(4-t-butylbenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (0.60 g, 56%); MS(FD): 489.2 (MH⁺).

Analysis for C₂₇H₂₇N₄O₅:


20 Calculated: C, 66.38; H, 5.78; N, 11.47
Found: C, 66.10; H, 5.80; N, 11.19

- 84 -

Example 26

Preparation of 4-methoxycarbonyl-N¹-[3-[(ethoxycarbonyl-amino)iminomethyl]benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

5




To a solution of 4-methoxycarbonyl-N¹-[4-(aminoimino-methyl)benzoyl]]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride (250 mg, 0.49 mmol) and K₂CO₃ (200 mg, 1.5 mmol) in THF (10 mL) and water (10 mL) was added a solution of ethyl chloroformate (0.07 mL, 0.74 mmol) in THF (5 mL). After 1 h, the reaction was concentrated and the residue was dissolved in ethyl acetate. The organic solution was washed with satd aq NH₄Cl (2X), water, satd aq NaHCO₃, water, brine, MgSO₄ dried, and concentrated. The residue was chromatographed (50% ethyl acetate/hexanes to 75% ethyl acetate/hexanes) to give the title compound as a solid (150 mg, 56%); FD-MS(FD): 544.

20

Example 27

Preparation of 4-methoxycarbonyl-N¹-[3-[(acetoxymethoxycarbonyl)amino]iminomethyl]benzoyl]-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

- 85 -




Using a procedure similar to Example 26 except starting
 5 from 4-methoxycarbonyl-N¹-[4-(aminoiminomethyl)benzoyl]-N²-
 (4-t-butylbenzoyl)-1,2-benzenediamine hydrochloride and
 acetoxymethyl 4-nitrophenyl carbonate, the title compound
 was obtained as a solid (54%); MS(FD): 589.

Analysis for C₃₁H₃₂N₄O₈:

10 Calculated: C, 63.26; H, 5.48; N, 9.52
 Found: C, 63.13; H, 5.57; N, 9.32

Example 28

Preparation of N¹-[3-[(methoxycarbonylamino)iminomethyl]-
 15 benzoyl]]-N²-(4-methoxybenzoyl)-1,2-benzenediamine



Using a procedure similar to Example 26 except starting
 20 from methyl chloroformate and N¹-[3-(aminoiminomethyl)-
 benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine
 hydrochloride, the title compound was obtained as a solid
 (74%); MS(FD): 446.

Analysis for C₂₄H₂₂N₄O₅:


- 86 -

Calculated: C, 64.56; H, 4.96; N, 12.55

Found: C, 64.43; H, 5.16; N, 12.07

Example 29

- 5 Preparation of N¹-[3-[(ethoxycarbonylamino)iminomethyl]benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine




- 10 Using a procedure similar to Example 26 except starting from N¹-[3-(aminoiminomethyl)benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride, the title compound was obtained as a solid (57%); MS(FD): 460.1.

Analysis for C₂₅H₂₄N₄O₅·0.9H₂O:

- 15 Calculated: C, 62.99; H, 5.46; N, 11.75
Found: C, 63.10; H, 5.33; N, 11.63

Example 30

- 20 Preparation of N¹-[3-[(1,1-dimethylethoxycarbonylamino)iminomethyl]benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine




- 87 -

Using a procedure similar to Example 26 except starting from N¹-[3-(aminoiminomethyl)benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride, the title compound was obtained as a solid (43%); MS(FD): 489.

- 5 Analysis for C₂₇H₂₈N₄O₅:
 Calculated: C, 66.38; H, 5.78; N, 11.47
 Found: C, 66.17; H, 5.79; N, 11.33

Example 31

- 10 Preparation of N¹-[2-hydroxy-3-(aminoiminomethyl)benzoyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride



- 15 A) N¹-(2-fluoro-3-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine

 Using a procedure similar to Example 1B except starting from 2-fluoro-3-cyanobenzoic acid and N¹-(4-methoxybenzoyl)-N²-benzenediamine, the title compound was obtained as a

- 20 solid (74%); MS(FD): 389.1.

- B) N¹-[(3-amino-1,2-benzisoxazol-7-yl)carbonyl]-N²-(4-methoxybenzoyl)-1,2-benzenediamine

- To a suspension of potassium t-butoxide (92 mg, 0.82
25 mmol) in DMF (5 mL) was added acetone oxime hydrochloride (60 mg, 0.82 mmol). After 1 h, N¹-(2-fluoro-3-cyanobenzoyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine (290 mg, 0.74 mmol) was added. After another 2 h, potassium
 t-butoxide (92 mg, 0.82 mmol) and acetone oxime
30 hydrochloride (60 mg, 0.82 mmol) were added. After 12 h,

- 88 -

the reaction was concentrated and the residue was dissolved in ethyl acetate and washed with water (2X), brine, MgSO_4 dried, and concentrated. The resulting solid was dissolved in ethanol (35 mL), 5% HCl (35 mL) was added, and the mixture was refluxed for 1 h. The reaction was cooled, concentrated to 2 mL, diluted with ethyl acetate (250 mL), and neutralized with satd aq NaHCO_3 . The organic layer was washed with satd aq NaHCO_3 (2X) and brine, MgSO_4 dried, concentrated and crystallized to give the title compound as an off-white solid (106 mg, 35%); MS(FD): 402.1.

Analysis for $\text{C}_{22}\text{H}_{18}\text{N}_4\text{O}_4$:

Calculated: C, 65.66; H, 4.51; N, 13.92

Found: C, 65.41; H, 4.61; N, 13.73

C) N^1 -[2-hydroxy-3-(aminoiminomethyl)benzoyl]- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 1C, except starting from N^1 -[(3-amino-1,2-benzisoxazol-7-yl)carbonyl]- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine, the title compound was obtained as a solid (20 mg, 32%); MS(ES): 405.1 (MH^+).

Analysis for $\text{C}_{22}\text{H}_{20}\text{N}_4\text{O}_4 \cdot 1.1\text{HCl} \cdot 1.4\text{H}_2\text{O}$:


Calculated: C, 56.25; H, 5.13; N, 11.92; Cl, 8.30

Found: C, 56.39; H, 4.63; N, 11.66; Cl, 8.38

25

Example 32

Preparation of N-[3-(aminoiminomethyl)benzoyl]-2-[(4-methoxyphenoxy)methyl]aniline hydrochloride



30

- 89 -

A) 2-[(4-methoxyphenoxy)methyl]nitrobenzene

To a suspension of potassium t-butoxide (5.7 g, 51 mmol) in THF (200 mL) was added 4-methoxyphenol (6.3 g, 51 mmol). After 10 min, 2-nitrobenzyl bromide (10 g, 46 mmol) was added. After 48 h, the reaction was concentrated and the residue was dissolved in ethyl acetate, washed with 1 N citric acid (2X), brine, satd aq NaHCO₃ (2X), water, and brine. The organic layer was MgSO₄ dried, concentrated, triturated with ether, filtered and vacuum dried to give the title compound as an off-white solid (8.4 g, 71%); MS(FD): 259.

Analysis for C₁₄H₁₃NO₄:

Calculated: C, 64.86; H, 5.05; N, 5.40

Found: C, 64.62; H, 5.05; N, 5.43

B) 2-[(4-methoxyphenoxy)methyl]aniline

Using a procedure similar to Example 1C except starting from 2-[(4-methoxyphenoxy)methyl]nitrobenzene, the title compound was obtained as a solid (2.0 g, 59%); MS(FD): 229.1.

Analysis for C₁₄H₁₅NO₂:

Calculated: C, 73.34; H, 6.59; N, 6.11

Found: C, 73.38; H, 6.46; N, 6.19

C) N-(3-cyanobenzoyl)-2-[(4-methoxyphenoxy)methyl]aniline

Using a procedure similar to Example 1B except starting from 2-[(4-methoxyphenoxy)methyl]aniline and 3-cyanobenzoyl chloride, the title compound was obtained as a solid (0.50 g, 36%); MS(FD): 359.

Analysis for C₂₂H₁₈N₂O₃:

Calculated: C, 73.73; H, 5.06; N, 7.82

Found: C, 73.99; H, 5.21; N, 7.87

- 90 -

D) N-[3-(aminoiminomethyl)benzoyl]-2-[(4-methoxyphenoxy)-methyl]-aniline hydrochloride

Using a procedure similar to Example 15 except starting from 1-[N-(3-cyanobenzoyl)]-2-[(4-methoxyphenyloxy)methyl]-

5 aniline, the title compound was obtained as a solid (18%);

MS(ES): 376.1 (MH⁺).

Analysis for C₂₂H₂₁N₃O₃·1.1HCl·0.3H₂O:


Calculated: C, 62.78; H, 5.44; N, 9.98; Cl, 9.26

Found: C, 62.99; H, 5.23; N, 10.05; Cl, 9.21

10

Example 33

Preparation of N¹-[3-(aminoiminomethyl)benzoyl]-N²-[(4-methoxybenzyl)oxy]-aniline hydrochloride



15

A) N-(3-cyanobenzoyl)-2-hydroxyaniline

Using a procedure similar to Example 1B except starting from 2-hydroxyaniline, the title compound was obtained as a solid (70%); MS(FD): 238.

20

Analysis for C₁₄H₁₀N₂O₂:

Calculated: C, 70.58; H, 4.23; N, 11.76

Found: C, 70.83; H, 3.99; N, 11.46

25 B) N-(3-cyanobenzoyl)-2-[(4-methoxybenzyl)oxy]aniline

Using a procedure similar to Example 32A except starting from 4-methoxybenzyl chloride and N¹-(3-cyanobenzoyl)-2-hydroxyaniline and using K₂CO₃/DMF in place of potassium t-butoxide/THF, the title compound was obtained as a solid (27%); MS(FD): 359.

30

- 91 -

Analysis for $C_{22}H_{18}N_2O_3$:

Calculated: C, 73.73; H, 5.06; N, 7.82

Found: C, 73.44; H, 4.98; N, 7.64

- 5 C) N-[3-(aminoiminomethyl)benzoyl]-2-[(4-methoxybenzyl)oxy]aniline hydrochloride

Using a procedure similar to Example 15, except starting from N^1 -(3-cyanobenzoyl)- N^2 -[(4-methoxybenzyl)oxy]aniline, the title compound was obtained as a solid (10%);

- 10 MS(ES): 376.1 (MH^+).

Analysis for $C_{22}H_{21}N_3O_3 \cdot 1.2HCl \cdot H_2O$:


Calculated: C, 60.44; H, 5.58; N, 9.61; Cl, 9.73

Found: C, 60.22; H, 5.22; N, 9.91; Cl, 9.96

15

Example 34

Preparation of 2-[3-(aminoiminomethyl)benzyloxy]-N-(4-methoxybenzoyl)aniline hydrochloride



20

- A) 2-(3-cyanobenzyloxy)-N-(4-methoxybenzoyl)aniline

Using a procedure similar to Example 33A and 33B except starting from 2-hydroxyaniline, p-anisoyl chloride and 3-cyanobenzyl bromide, the title compound was obtained as a solid (700 mg, 48%); MS(FD): 358.1.

25

Analysis for $C_{22}H_{18}N_2O_3$:

Calculated: C, 73.73; H, 5.06; N, 7.82

Found: C, 73.67; H, 5.27; N, 7.61

- 92 -

B) 2-[3-(aminoiminomethyl)benzyloxy]-N-(4-methoxybenzoyl)-aniline hydrochloride

Using a procedure similar to Example 15, except starting from 2-(3-cyanobenzyloxy)-N-(4-methoxybenzoyl)-aniline, the title compound was obtained as a solid (10%); MS(ES): 376.2 (MH⁺).


Analysis for C₂₂H₂₁N₃O₃·1.1HCl·0.5H₂O:

Calculated: C, 62.24; H, 5.48; N, 9.89; Cl, 9.19

Found: C, 62.33; H, 5.32; N, 9.97; Cl, 9.02

Example 35

Preparation of N¹-[[1-(amino(imino)methyl))piperidin-3-yl]-carbonyl]-N²-(t-butylbenzoyl)-1,2-benzenediamine trifluoroacetate



A) 1-(benzyloxycarbonylamino(benzyloxycarbonylimino)-methyl)piperidine-3-carboxylic acid

To a solution of nipecotic acid (1.66 g, 12.8 mmol) in 2 N NaOH (12.8 mL, 25.6 mmol) and dioxane (30 mL) was added 1,3-bis(benzyloxycarbonyl)-2-methyl-2-thiopseudourea (4.6 g, 12.8 mmol). After 48 h, the reaction mixture was added to H₂O (200 mL) and Et₂O (200 mL). The aqueous layer was separated and extracted with Et₂O (2 X 200 mL). The aqueous layer was acidified to pH 2.0 with 3 N HCl and extracted with EtOAc (500 mL). The organic layer was MgSO₄ dried and

- 93 -

concentrated to give the title compound (4.0 g, 71%): TLC
R_f=0.65 (1:15: 135 Acetic Acid:Methanol:Chloroform); NMR
(CDCl₃) δ 1.55-1.92 (m, 6H), 2.00-2.15 (m, 1H), 2.70-2.80
(m, 1H), 3.20-3.35 (q, 1H), 3.38-3.45 (m, 1H), 3.75-4.10 (m,
5 1H), 5.10 (s, 2H), 5.22 (s, 2H), 7.30-7.40 (m, 10H); FD-MS
440 (MH⁺).

Analysis for C₂₃H₂₅N₃O₆:

Calculated: C, 62.86; H, 5.73; N, 9.56

Found: C, 62.64, H, 5.79, N, 9.38

10

B) N¹-[[1-(benzyloxycarbonylamino(benzyloxycarbonylimino)-
methyl)piperidin-3-yl]carbonyl]-N²-(t-butylbenzoyl)-1,2-
benzenediamine

To a solution of 1-(benzyloxycarbonylamino(benzyloxy-
15 carbonylamino)methyl)piperidine-3-carboxylic acid (1.2 g,
2.7 mmol) in CH₂Cl₂ (30 mL) was added 2-(4-t-butylbenzoyl-
amino)aniline (obtained from 4-t-butylbenzoyl chloride and
2-nitroaniline using procedures similar to those of Example
1-B and 1-C) (0.730 g, 2.7 mmol), DCC (0.556 g, 2.7 mmol),
20 and HOAt (0.367 g, 2.7 mmol). The reaction was stirred for
24 h. The resultant precipitate was filtered and the
filtrate was concentrated. The residue was dissolved in
EtOAc (250ml) and washed with 1N NaHCO₃, water, and 1.5 N
citric acid. The organic layer was MgSO₄ dried and
25 concentrated to give the title compound as an amorphous
solid (2.4 g, 100%).

C) N¹-[[1-(amino(imino)methyl)piperidin-3-yl]carbonyl]-
N²-(t-butylbenzoyl)-1,2-benzenediamine trifluoroacetate

30 A mixture of N¹-[[1-(benzyloxycarbonylamino(benzyloxy-
carbonylimino)methyl)piperidin-3-yl]carbonyl]-N²-(t-butyl-
benzoyl)-1,2-benzenediamine (2.4 g), ethanol (60 mL), water
(25 mL), 1 N HCl (5.0 mL, 5.0 mmol), and 5% Pd / C catalyst
(1.5 g) was hydrogenated at 1 atm. After the reaction was

- 94 -

completed, the catalyst was filtered through diatomaceous earth. The filtrate was concentrated and vacuum dried overnight. The residue was dissolved in 0.1% TFA, filtered through a Millipore 0.5 mm filter, and chromatographed on a 5 X 25 cm column Vydac C18 resin (5% CH₃CN/H₂O to 50% CH₃CN/H₂O) to give the title compound as a white solid (0.550 g, 37%); MS(FAB): 422 (MH⁺).


Analysis for C₂₆H₃₅F₃N₅O₄·H₂O:

Calculated: C, 56.41; H, 5.83; N, 12.65

Found: C, 56.41, H, 5.94, N, 12.27.

Example 36

Preparation of N¹-[[1-(amino(imino)methyl)piperidin-3-yl]-carbonyl]-N²-(4-t-butylbenzoyl)-4-methoxycarbonyl-1,2-benzenediamine hydrochloride



A) 4-methoxycarbonyl-N²-(4-t-butylbenzoyl)-1,2-benzenediamine

Using a procedure similar to Example 19A except starting with 4-tert-butylbenzoyl chloride and crystallization of the crude product from 1:1:1 CH₂Cl₂:Et₂O:hexane, gave the title compound as a solid (76%); TLC: R_f=0.65 (ethyl acetate:hexane 1:1); MS(FAB): 327 (MH⁺).

Analysis for C₁₉H₂₂N₂O₃:

Calculated: C, 69.92; H, 6.79; N, 8.58

Found: C, 69.69, H, 6.63, N, 8.5

- 95 -

B) N¹-[[1-(benzyloxycarbonylamino(benzyloxycarbonylimino)-methyl)piperidin-3-yl]carbonyl]-N²-(4-t-butylbenzoyl)-4-methoxycarbonyl-1,2-benzenediamine

Using a procedure similar to Example 35B except
5 starting with 4-methoxycarbonyl-N²-(4-t-butylbenzoyl)-1,2-benzenediamine and using chromatography (CHCl₃ to 30% EtOAc/CHCl₃), the title compound was obtained as an amorphous solid (1.54 g, .44%); TLC: R_f=0.29 (8:2 chloroform:ethyl acetate); MS(FD): 748 (MH⁺).

10 Analysis for C₄₂H₄₅N₅O₈:

Calculated: C, 67.46; H, 6.07; N, 9.36

Found: C, 67.47, H, 5.98, N, 9.10

C) N¹-[[1-(amino(imino)methyl)piperidin-3-yl]carbonyl]-
15 N²-(4-t-butylbenzoyl)-4-methoxycarbonyl-1,2-benzenediamine hydrochloride

Using a procedure similar to Example 35 except starting with N¹-[[1-(benzyloxycarbonylamino(benzyloxycarbonylimino)-methyl)piperidin-3-yl]carbonyl]-N²-(4-t-butylbenzoyl)-4-methoxycarbonyl-1,2-benzenediamine hydrochloride followed by
20 precipitation with ether, the title compound was obtained as a white solid (78%); MS(FAB): 480 (MH⁺).

Analysis for C₂₆H₃₃N₅O₄·HCl·2.5H₂O:


Calculated: C, 55.65; H, 6.83; N, 12.48

25 Found: C, 55.76, H, 6.75, N, 12.34

Example 37

Preparation of N¹-[[1-(amino(imino)methyl)piperidin-3-yl]carbonyl]-N²-(4-t-butylbenzoyl)-4-carboxy-1,2-
30 benzenediamine hydrochloride

- 96 -



A) N¹-[[1-(benzyloxycarbonylamino(benzyloxycarbonylimino)-
methyl)piperidin-3-yl]carbonyl]-N²-(4-t-butylbenzoyl)-4-
5 carboxy-1,2-benzenediamine

To a solution of N¹-[[1-(benzyloxycarbonylamino-
(benzyloxycarbonylimino)methyl)piperidin-3-yl]carbonyl]-N²-
(4-t-butylbenzoyl)-4-methoxycarbonyl-1,2-benzenediamine (0.7
g, 0.94 mmol), CH₃OH (10 mL), dioxane (10 mL), and water
10 (6 mL) was added 1 N NaOH (3.7 mL, 3.7 mmol). The reaction
was stirred for 48 h and was concentrated. The residue was
partitioned between H₂O (100 mL) and Et₂O (100 mL). The
aqueous layer was extracted with Et₂O (100 mL), acidified to
pH 2.5 with 5 N HCl and extracted with EtOAc (3 X 200 mL).
15 The combined EtOAc layers were washed with water, MgSO₄
dried, concentrated and crystallized at 0 °C overnight to
give the title compound as a solid (0.63 g, 92%); TLC:
R_f=0.38 (1:15: 135 acetic acid:methanol:chloroform); NMR
(CDCl₃) δ 1.35 (s, 9H), 1.50-2.05 (m, 8H), 3.20-3.40 (m, 1H),
20 3.60-4.05 (m, 3H), 5.05 (s, 4H), 7.20-7.35 (m, 12H), 7.40-
7.50 (m, 2H), 7.55-7.65 (m, 1H), 7.90-8.00 (m, 2H), 8.05-
8.15 (m, 1H); MS(FAB): 734 (MH⁺).

B) N¹-[[1-(amino(imino)methyl)piperidin-3-yl]carbonyl]-
25 N²-(4-t-butylbenzoyl)-4-carboxy-1,2-benzenediamine
hydrochloride

Using a procedure similar to Example 35, except
starting with N¹-[[1-(benzyloxycarbonylamino(benzyloxy-

- 97 -

carbonylimino)methyl)piperidin-3-yl]carbonyl]-N²-(4-t-butylbenzoyl)-4-carboxy-1,2-benzenediamine, the title compound was obtained as a white solid (79%); MS(FAB): 466 (MH⁺).

Analysis for C₂₅H₃₁N₅O₄·HCl·1.5H₂O:

- 5 Calculated: C, 56.76; H, 6.67; N, 13.24
 Found: C, 56.84; H, 6.29; N, 13.52

Example 38

Preparation of N¹-[3-(amino(imino)methyl)benzoyl]-N²-

10 (4-ethoxybenzoyl)-1,2-benzenediamine

A) N¹-(3-cyanobenzoyl)-1,2-benzenediamine

- Using a procedure similar to 1B except starting with 1,2-phenylenediamine and triethylamine as base, the title
15 compound was obtained as a white solid (45%); mp: 197-
200 °C; NMR(300 MHz, DMSO-d₆): δ 4.98 (s, 2H), 6.56 (t, 1H),
6.74 (d, 1H), 6.96 (t, 1H), 7.13 (d, 1H), 7.71 (t, 1H), 8.02
(d, 1H), 8.25 (d, 1H), 8.42 (s, 1H), 9.80 (s, 1H); MS(FD):
237.

20 Analysis for C₁₄H₁₁N₃O:

- Calculated: C, 70.87; H, 4.67; N, 17.71
 Found: C, 70.68; H, 4.58; N, 17.52

B) N¹-(3-cyanobenzoyl)-N²-(4-ethoxybenzoyl)-1,2-benzene-
25 diamine

- Using a procedure similar to 1B except starting with
N¹-(3-cyanobenzoyl)-1,2-benzenediamine, 4-ethoxybenzoyl
chloride and triethylamine as base, the title compound was
obtained as a white solid (39%); mp: 168-9 °C; IR(KBr):
30 1645, 1672, 2231, 3377 cm⁻¹; MS(FD): 385(M⁺).
NMR(300 MHz, DMSO-d₆): δ 1.31 (t, 3 H); 4.07 (quartet, 2 H);
7.03 (d, 2 H); 7.30 (m, 2 H); 7.60 (m, 2 H); 7.72 (t, 1 H);
7.91 (d, 2 H); 8.04 (d, 2 H); 8.21 (d, 1 H); 8.35 (s, 1 H)

- 98 -

Analysis for C₂₃H₁₉N₃O₃

Calculated: C, 71.68; H, 4.97; N, 10.90

Found: C, 69.54; H, 4.71; N, 10.65

- 5 C) N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-ethoxybenzoyl)-1,2-benzenediamine

Using a procedure similar to 1F except starting with N¹-(3-cyanobenzoyl)-N²-(4-ethoxybenzoyl)-1,2-benzenediamine, the title compound was obtained (100%) as a white solid;

- 10 mp 180-182 °C; NMR(300 MHz, DMSO-d₆): δ 1.31 (t, 3 H); 4.07 (quartet, 2 H); 7.01 (d, 2 H); 7.30 (m, 2 H); 7.50 (t, 1 H); 7.60 (m, 2 H); 7.84 (d, 1 H); 7.90 (m, 3 H); 8.25 (s, 1 H); 9.73 (s, 1 H); 9.89 (s, 1 H); 10.12 (s, 1 H)

Analysis for C₂₃H₂₂N₄O₄

- 15 Calculated: C, 66.02; H, 5.30; N, 13.39

Found: C, 65.39; H, 5.73; N, 12.63

D) N¹-[3-(amino(imino)methyl)benzoyl]-N²-(4-ethoxybenzoyl)-1,2-benzenediamine

- 20 Using a procedure similar to 1G except starting with N¹-[3-(amino(hydroxyimino)methyl)benzoyl]-N²-(4-ethoxybenzoyl)-1,2-benzenediamine, the title compound was obtained as a white solid (21%); mp 163-5 °C; NMR(300 MHz, DMSO-d₆): 1.31 (t, 3 H); 4.07 (quintet, 2 H); 7.01 (d, 2 H); 7.20 (m, 2 H); 7.60 (m, 2 H); 7.65 (t, 1 H); 8.00 (m, 3 H); 8.29 (d, 1 H); 8.54 (s, 1 H); 9.33 (s, 1 H); 9.54 (s, 1 H); 10.14 (s, 1 H); 10.72 (s, 1 H); MS(FD): 402.

Calculated: C, 62.94; H, 5.28; N, 12.76


Found: C, 45.54; H, 3.85; N, 8.92

30

Example 39

Preparation of N¹-(3-Amino-1,2-benzoxazol-5-ylcarbonyl)-N²-(4-methoxybenzoyl)-1,2-benzenediamine

- 99 -



A) N^1 -(3-Bromo-4-fluorobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine

5 By methods substantially equivalent to those described in Example 18, except without the use of N,N-diisopropylethylamine, N^1 -(3-bromo-4-fluorobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine (2.4 g, 69%) was prepared from N^1 -(4-methoxybenzoyl)-1,2-benzenediamine and 3-bromo-4-fluorobenzoic acid.

^1H NMR

FAB-MS, m/e 443.1 (MH^+)

Anal. for $\text{C}_{21}\text{H}_{16}\text{BrFN}_2\text{O}_3$

15 Calc: C, 56.90; H, 3.64; N, 6.32;
Found: C, 56.98; H, 3.55; N, 6.40.

B) N^1 -(3-Cyano-4-fluorobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine

20 CuCN (606 mg, 6.77 mmol) was dissolved in NMP (25 mL) and the solution was heated to 200 °C for 45 min. To this stirring solution was added N^1 -(3-bromo-4-fluorobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine (700 mg, 2.26 mmol) and stirring continued for 20 h. The solution was then cooled to room temperature, diluted with ethyl acetate and washed once with 1 N ammonium hydroxide, twice with 1 N HCl and once with brine. The organic phase was then dried with MgSO_4 , filtered and concentrated in vacuo. The residue was chromatographed with 10% ethyl acetate/dichloromethane to give 270 mg (44%) of off-white solid.

30 ^1H NMR

- 100 -

FD-MS, m/e 389.1 (M^+)Anal. for $C_{22}H_{16}N_3O_3 \cdot 0.9H_2O$

Calc: C, 68.35; H, 4.64; N, 10.86;

Found: C, 68.63; H, 4.58; N, 10.16.

5

C) N^1 -(3-Amino-1,2-benzoxazol-5-ylcarbonyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine

To a stirring solution of acetone oxime (94 mg, 1.28 mmol) in DMF (2 mL) was added a 1 M solution of potassium t-butoxide (1.28 mL, 1.28 mmol) in THF. After 30 min, N^1 -(3-cyano-4-fluorobenzoyl)- N^2 -(4-methoxybenzoyl)-1,2-benzenediamine (100 mg, 0.26 mmol) was added and stirring continued for 72 h. The solvents were removed in vacuo and the residue was dissolved in methanol (20 mL). To this stirring solution was added water (15 mL) and conc HCl (3 mL), and the solution was heated to reflux. After 2 h, the solvents were removed in vacuo and the residue was dissolved in methanol. Silica gel was added and then the solvents were removed in vacuo. This dry pack was loaded onto a silica gel column and eluted with 1:1 ethyl acetate/dichloromethane. The product containing fractions were combined and concentrated to give 48 mg (46%) of white solid.

1H NMR

25 FD-MS, m/e 402.1 (M^+)Anal. for $C_{22}H_{18}N_4O_4 \cdot 0.9H_2O$

Calc: C, 63.13; H, 4.77; N, 13.38;

Found: C, 63.24; H, 4.52; N, 12.31.

30 Using procedures analogous to those above and standard procedures of organic chemistry, also were prepared the following compounds:

4-Acetylamino- N^1 -[3-(amino(imino)methyl)benzoyl]- N^2 -(4-t-butylbenzoyl)-1,2-benzenediamine;


- 101 -

4-Acetylamino-N¹-[3-(amino(imino)methyl)benzoyl]-N²-
[1-(4-pyridyl)piperidin-4-yl]methoxycarbonyl-1,2-
benzenediamine.

- 102 -

What is claimed is:

1. A method of inhibiting factor Xa comprising using
an effective amount of a factor Xa inhibiting compound of
5 formula I



wherein

- A³, A⁴, A⁵ and A⁶, together with the two carbons to
10 which they are attached, complete a substituted benzene in
which A³ is CR³, A⁴ is CR⁴, A⁵ is CR⁵, and A⁶ is CR⁶;

wherein

- R³ is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy
(which may bear an ω-carboxy substituent), benzyloxy (which
15 may bear one or more halo, hydroxy methoxy or methyl
substitutents), methyl or methoxy;

- one of R⁴ and R⁵ is hydrogen, methyl, halo, trifluoro-
methyl, nitro, amino(imino)methyl, amino(hydroxyimino)-
methyl, R^fO-, R^fO₂C-, R^fO₂C-CH₂-, R^fO₂C-CH₂-O-,
20 3-methoxycarbonyl-1-oxopropyl, R^gNH- or bis(methylsulfonyl)-
amino;

- the other of R⁴ and R⁵ is hydrogen, halo or methyl; and
R⁶ is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy
(which may bear an ω-carboxy substituent), benzyloxy (which
25 may bear one or more halo, hydroxy methoxy or methyl
substitutents), methyl or methoxy;

- in which R^f is hydrogen, (1-4C)alkyl or benzyl; R^g is
hydrogen, acetyl, trifluoroacetyl, phenylalanyl,
2-(t-butoxycarbonylamino)-4-methylsulfinyl-1-oxobutyl or
30 R^hSO₂-; and R^h is (1-4C)alkyl, trifluoromethyl, phenyl,
3,5-dimethylisoxazol-4-yl or dimethylamino; or

- 103 -

two adjacent residues selected from R^3 , R^4 , R^5 and R^6 together form a benz ring; and the other two are each hydrogen; or

A^3 , A^4 , A^5 and A^6 , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which

(a) one of A^3 , A^4 , A^5 and A^6 is N, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively;

(b) two adjacent residues of A^3 , A^4 , A^5 and A^6 together form S, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively;

(c) two non-adjacent residues of A^3 , A^4 , A^5 and A^6 are each N, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively; or

(d) A^3 and A^4 together form a fused benz ring, and A^5 and A^6 together form -NH-;


wherein

each of R^3 , R^4 , R^5 and R^6 is hydrogen, or one or two of R^3 , R^4 , R^5 and R^6 is independently chloro, bromo or methyl and the others are hydrogen;

L^1 is -NH-CO-, -CO-NH- or -O-CH₂- such that - L^1 - Q^1 is -NH-CO- Q^1 , -CO-NH- Q^1 or -O-CH₂- Q^1 (provided that when - L^1 - Q^1 is -CO-NH- Q^1 , then R^2 is not -CO-NH- Q^1 or -CO-NH- Q^{2B});


Q^1 is Q^{1A} , Q^{1B} or Q^{1C} wherein

Q^{1A} is




in which R^a is hydrogen, R^b is hydrogen and R^c is hydrogen or hydroxy; or R^a is hydroxy, R^b is hydrogen and R^c is hydrogen; or R^a and R^b together form a methylene or oxo group and R^c is hydrogen;


- 104 -

Q^{1B} is

in which R^d is hydrogen or hydroxy and R^e is hydrogen or
 5 fluoro; and

Q^{1C} is


R² is a residue defined for -NH-CO-Q¹ or -CO-NH-Q¹; or
 10 R² is -L^{2A}-Q^{2A}, -L^{2B}-Q^{2B}, -L^{2C}-Q^{2C} or -L^{2D}-Q^{2D} wherein
 L^{2A} is a direct bond; and
 Q^{2A} is



15 in which D is carbonyl or -CHR^k- in which R^k is hydrogen,
 hydroxy, (1-6C)alkoxy or -CH₂-R^j in which R^j is carboxy,
 [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or
 two (1-2C)alkyl substituents on the nitrogen; and one of R^m
 and Rⁿ is hydrogen and the other is amino, bromo,
 20 (1-4C)alkyl or (1-4C)alkoxy, or R^m and Rⁿ together form a
 benz ring;

L^{2B} is -NH-CO-, -CO-NH-, -O-CO-, -CH₂-O- or -O-CH₂-
 such that -L^{2B}-Q^{2B} is -NH-CO-Q^{2B}, -CO-NH-Q^{2B}, -O-CO-Q^{2B},
 -CH₂-O-Q^{2B} or -O-CH₂-Q^{2B}; and

- 105 -

Q^{2B} is

- in which R^O is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and R^P is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl, 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or -J-R^Q in which J is a single bond, methylene, carbonyl, oxo, -S(O)_q- (wherein q is 0, 1 or 2), or -NR^r- (wherein R^r is hydrogen or methyl); and R^Q is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;
- L^{2C} is -NR^V-CO-X-, -NR^V-CS-Y-, -CH₂-CO-NR^W-CH₂-, -O-CO-, -O-CH₂-, -S-CH₂- or -CH₂-NR^X-CH₂- such that -L^{2C}-Q^{2C} is -NR^V-CO-X-Q^{2C}, -NR^V-CS-Y-Q^{2C}, -CH₂-CO-NR^W-CH₂-Q^{2C}, -O-CO-Q^{2C}, -O-CH₂-Q^{2C}, -S-CH₂-Q^{2C} or -CH₂-NR^X-CH₂-Q^{2C} in which X is -(CH₂)_x- (wherein x is 0, 1 or 2), -NR^W-CH₂-, -O-CH₂- or -S-CH₂-; Y is -NR^W-CH₂- or -O-CH₂-; each of R^V and R^W is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the α-position; and R^X is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and
- Q^{2C} is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;
- L^{2D} is -NH-CO- such that -L^{2D}-Q^{2D} is -NH-CO-Q^{2D}; and
- Q^{2D} is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and -CH₂-R^Z in which R^Z is isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl

- 106 -

in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy, acetoxy, benzyloxy, amino, acetylamino, nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a
5 methyl or nitro substituent;

or a prodrug of the compound of formula I;

or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

10 2. The method of Claim 1 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl or ethyl; (1-4C)alkyl is methyl, ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl; (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or
15 hexyl; and halo is bromo or chloro.

3. The method of Claim 2 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl; (1-4C)alkyl is methyl, isopropyl,
20 butyl or t-butyl; (1-6C)alkyl is methyl, butyl or hexyl; and halo is chloro.

4. The method of any of the above Claims 1-3 wherein R^2 is selected from $-L^{2A}-Q^{2A}$, $-NH-CO-Q^{2B}$, $-NR^V-CO-X-Q^{2C}$,
25 $-NR^V-CS-Y-Q^{2C}$, and $-NH-CO-Q^{2D}$.

5. The method of any of the above Claims 1-4 wherein the compound of formula I is one in which each of A^3 , A^5 and A^6 is CH.

30

6. The method of any of the above Claims 1-5 wherein R^2 is (4-isopropylbenzoyl)amino, (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)amino, or [1-(4-pyridyl)piperidin-4-yl]-methoxycarbonylamino.

- 107 -


7. The method of any of the above Claims 1-6 wherein R^4 or R^5 is hydrogen, hydroxy, acetylamino, carboxymethoxy, carboxymethyl, methoxycarbonyl or carboxy.

5

8. The method of any of the above Claims 1-7 wherein L^1-Q^1 is $-NH-CO-Q^1$.

9. The method of any of the above Claims 1-7 wherein
10 L^1-Q^1 is $-CO-NH-Q^1$.

10. A novel compound of formula I



15 wherein

A^3 , A^4 , A^5 and A^6 , together with the two carbons to which they are attached, complete a substituted benzene in which A^3 is CR^3 , A^4 is CR^4 , A^5 is CR^5 , and A^6 is CR^6 ;
wherein

20 R^3 is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy (which may bear an ω -carboxy substituent), benzoyloxy (which may bear one or more halo, hydroxy methoxy or methyl substitutents), methyl or methoxy;

one of R^4 and R^5 is hydrogen, methyl, halo, trifluoro-
25 methyl, nitro, amino(imino)methyl, amino(hydroxyimino)-methyl, R^fO- , R^fO_2C- , $R^fO_2C-CH_2-$, $R^fO_2C-CH_2-O-$, 3-methoxycarbonyl-1-oxopropyl, R^9NH- or bis(methylsulfonyl)-amino;

the other of R^4 and R^5 is hydrogen, halo or methyl; and
30 R^6 is hydrogen, hydroxy, [(1-2C)alkyl]carbonyloxy (which may bear an ω -carboxy substituent), benzoyloxy (which

- 108 -

may bear one or more halo, hydroxy methoxy or methyl substitutents), methyl or methoxy;

in which R^f is hydrogen, (1-4C)alkyl or benzyl; R^g is hydrogen, acetyl, trifluoroacetyl, phenylalanyl,

5 2-(t-butoxycarbonylamino)-4-methylsulfinyl-1-oxobutyl or R^hSO_2- ; and R^h is (1-4C)alkyl, trifluoromethyl, phenyl, 3,5-dimethylisoxazol-4-yl or dimethylamino; or

two adjacent residues selected from R^3 , R^4 , R^5 and R^6 together form a benz ring; and the other two are each

10 hydrogen; or

A^3 , A^4 , A^5 and A^6 , together with the two carbons to which they are attached, complete a substituted heteroaromatic ring in which

(a) one of A^3 , A^4 , A^5 and A^6 is N, and each of the others
15 is CR^3 , CR^4 , CR^5 or CR^6 , respectively;

(b) two adjacent residues of A^3 , A^4 , A^5 and A^6 together form S, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively;

(c) two non-adjacent residues of A^3 , A^4 , A^5 and A^6 are each
20 N, and each of the others is CR^3 , CR^4 , CR^5 or CR^6 , respectively; or

(d) A^3 and A^4 together form a fused benz ring, and A^5 and A^6 together form -NH-;

wherein


25 each of R^3 , R^4 , R^5 and R^6 is hydrogen, or one or two of R^3 , R^4 , R^5 and R^6 is independently chloro, bromo or methyl and the others are hydrogen;

L^1 is -NH-CO-, -CO-NH- or -O-CH₂- such that - L^1 - Q^1 is -NH-CO- Q^1 , -CO-NH- Q^1 or -O-CH₂- Q^1 (provided that when - L^1 - Q^1
30 is -CO-NH- Q^1 , then R^2 is not -CO-NH- Q^1 or -CO-NH- Q^{2B});

Q^1 is Q^{1A} , Q^{1B} or Q^{1C} wherein


Q^{1A} is

- 109 -




- in which R^a is hydrogen, R^b is hydrogen and R^c is hydrogen or hydroxy; or R^a is hydroxy, R^b is hydrogen and R^c is hydrogen; or R^a and R^b together form a methylene or oxo group and R^c is hydrogen;

Q^{1B} is




- 10 in which R^d is hydrogen or hydroxy and R^e is hydrogen or fluoro; and

Q^{1C} is



- 15 R^2 is a residue defined for $-NH-CO-Q^1$ or $-CO-NH-Q^1$; or R^2 is $-L^{2A}-Q^{2A}$, $-L^{2B}-Q^{2B}$, $-L^{2C}-Q^{2C}$ or $-L^{2D}-Q^{2D}$ wherein L^{2A} is a direct bond; and Q^{2A} is




20

- 110 -

in which D is carbonyl or $-\text{CHR}^k-$ in which R^k is hydrogen, hydroxy, (1-6C)alkoxy or $-\text{CH}_2-\text{R}^j$ in which R^j is carboxy, [(1-4C)alkoxy]carbonyl or carbamoyl which may bear one or two (1-2C)alkyl substituents on the nitrogen; and one of R^m and R^n is hydrogen and the other is amino, bromo, (1-4C)alkyl or (1-4C)alkoxy, or R^m and R^n together form a benz ring;

$\text{L}^{2\text{B}}$ is $-\text{NH}-\text{CO}-$, $-\text{CO}-\text{NH}-$, $-\text{O}-\text{CO}-$, $-\text{CH}_2-\text{O}-$ or $-\text{O}-\text{CH}_2-$ such that $-\text{L}^{2\text{B}}-\text{Q}^{2\text{B}}$ is $-\text{NH}-\text{CO}-\text{Q}^{2\text{B}}$, $-\text{CO}-\text{NH}-\text{Q}^{2\text{B}}$, $-\text{O}-\text{CO}-\text{Q}^{2\text{B}}$, $-\text{CH}_2-\text{O}-\text{Q}^{2\text{B}}$ or $-\text{O}-\text{CH}_2-\text{Q}^{2\text{B}}$; and $\text{Q}^{2\text{B}}$ is



in which R^{O} is hydrogen, halo, (1-6C)alkyl, (1-4C)alkoxy, benzyloxy or (1-4C)alkylthio; and R^{P} is 1-hydroxyethyl, 1-hydroxy-1-methylethyl, 1-methoxy-1-methylethyl, 4-piperidinyl, 4-pyridinyl, dimethylaminosulfonyl or $-\text{J}-\text{R}^{\text{Q}}$ in which J is a single bond, methylene, carbonyl, oxo, $-\text{S}(\text{O})_{\text{q}}-$ (wherein q is 0, 1 or 2), or $-\text{NR}^{\text{F}}-$ (wherein R^{F} is hydrogen or methyl); and R^{Q} is (1-6C)alkyl, phenyl, 3-pyridyl or 4-pyridyl;

$\text{L}^{2\text{C}}$ is $-\text{NR}^{\text{V}}-\text{CO}-\text{X}-$, $-\text{NR}^{\text{V}}-\text{CS}-\text{Y}-$, $-\text{CH}_2-\text{CO}-\text{NR}^{\text{W}}-\text{CH}_2-$, $-\text{O}-\text{CO}-$, $-\text{O}-\text{CH}_2-$, $-\text{S}-\text{CH}_2-$ or $-\text{CH}_2-\text{NR}^{\text{X}}-\text{CH}_2-$ such that $-\text{L}^{2\text{C}}-\text{Q}^{2\text{C}}$ is $-\text{NR}^{\text{V}}-\text{CO}-\text{X}-\text{Q}^{2\text{C}}$, $-\text{NR}^{\text{V}}-\text{CS}-\text{Y}-\text{Q}^{2\text{C}}$, $-\text{CH}_2-\text{CO}-\text{NR}^{\text{W}}-\text{CH}_2-\text{Q}^{2\text{C}}$, $-\text{O}-\text{CO}-\text{Q}^{2\text{C}}$, $-\text{O}-\text{CH}_2-\text{Q}^{2\text{C}}$, $-\text{S}-\text{CH}_2-\text{Q}^{2\text{C}}$ or $-\text{CH}_2-\text{NR}^{\text{X}}-\text{CH}_2-\text{Q}^{2\text{C}}$ in which X is $-(\text{CH}_2)_x-$ (wherein x is 0, 1 or 2), $-\text{NR}^{\text{W}}-\text{CH}_2-$, $-\text{O}-\text{CH}_2-$ or $-\text{S}-\text{CH}_2-$; Y is $-\text{NR}^{\text{W}}-\text{CH}_2-$ or $-\text{O}-\text{CH}_2-$; each of R^{V} and R^{W} is independently hydrogen, benzyl or (1-6C)alkyl which is not branched at the α -position; and R^{X} is hydrogen, benzyloxycarbonyl or [(1-4C)alkoxy]carbonyl; and

- 111 -

Q^{2C} is 1-(4-pyridyl)piperidin-4-yl in which the pyridyl may bear a substituent at its 2-position selected from cyano, aminomethyl, carboxy, hydroxymethyl and (1-2C)alkyl;

L^{2D} is -NH-CO- such that -L^{2D}-Q^{2D} is -NH-CO-Q^{2D}; and

5 Q^{2D} is selected from 4-(4-pyridinyl)benzyloxy, 9-oxo-9H-fluoren-3-yl, benzo[b]thiophen-2-yl (which may bear a chloro, methyl or methoxy substituent), benzofuran-2-yl (which may bear a chloro, methyl or methoxy substituent), 4-(4-morpholinyl)-4-oxobutyl, and 4-piperidinyl bearing a
10 substituent at the 1-position selected from methylsulfonyl, phenylsulfonyl and -CH₂-R^Z in which R^Z is isopropyl, cyclopropyl, phenyl, furyl, thienyl, 2-thiazolyl, or pyridyl in which the phenyl may bear one or two substituents independently selected from halo, cyano, hydroxy, methoxy,
15 acetoxyl, benzyloxy, amino, acetylamino, nitro and 3,4-methylenedioxy, and the thienyl or furyl may bear a methyl or nitro substituent;

or a prodrug of the compound of formula I;

20 or a pharmaceutically acceptable salt of the compound of formula I or prodrug thereof.

11. The compound of Claim 10 wherein for an alkyl group or the alkyl portion of an alkyl containing group, (1-2C)alkyl is methyl or ethyl; (1-4C)alkyl is methyl,
25 ethyl, propyl, isopropyl, butyl, isobutyl, or t-butyl; (1-6C)alkyl is methyl, ethyl, propyl, butyl, pentyl or hexyl; and halo is bromo or chloro.

12. The compound of Claim 11 wherein for an alkyl group or the alkyl portion of an alkyl containing group,
30 (1-2C)alkyl is methyl; (1-4C)alkyl is methyl, isopropyl, butyl or t-butyl; (1-6C)alkyl is methyl, butyl or hexyl; and halo is chloro.

- 112 -

13. The compound of any of the above Claims 10-12 wherein R^2 is selected from $-L^{2A}-Q^{2A}$, $-NH-CO-Q^{2B}$, $-NR^V-CO-X-Q^{2C}$, $-NR^V-CS-Y-Q^{2C}$, and $-NH-CO-Q^{2D}$.

5 14. The compound of any of the above Claims 10-13 wherein the compound of formula I is one in which each of A^3 , A^5 and A^6 is CH.

10 15. The compound of any of the above Claims 10-14 wherein R^2 is (4-isopropylbenzoyl)amino, (4-t-butylbenzoyl)amino, (4-methoxybenzoyl)amino, or [1-(4-pyridyl)-piperidin-4-yl]methoxycarbonylamino.

15 16. The compound of any of the above Claims 10-15 wherein R^4 or R^5 is hydrogen, hydroxy, acetylamino, carboxymethoxy, carboxymethyl, methoxycarbonyl or carboxy.

20 17. The compound of any of the above Claims 10-16 wherein L^1-Q^1 is $-NH-CO-Q^1$.

18. The compound of any of the above Claims 10-16 wherein L^1-Q^1 is $-CO-NH-Q^1$.


25 19. A prodrug of Claim 10 which is a carbamate in which an amino (or imino) group of Q^1 is substituted by a [(1-4C)alkoxy]-carbonyl or acetoxymethoxycarbonyl group.

30 20. A pharmaceutical composition comprising a compound of formula I, or prodrug or pharmaceutically acceptable salt thereof, as claimed in Claim 10 in association with a pharmaceutically acceptable carrier, excipient or diluent.

- 113 -


21. A process for preparing a compound of formula I (or a pharmaceutically acceptable salt thereof) as provided in Claim 10 which comprises

- (A) for a compound of formula I in which the linkage of R² to the ring terminates in -NH-CO-, -NR^V-CO- or -NR^V-CS-, acylating an amine of formula II,




- or a corresponding amine in which the nitrogen bears the group R^V, using a corresponding acid which terminates with the group HO-CO- or HO-CS-, or an activated derivative thereof;

- (B) for a compound of formula I in which -L¹-Q¹ is -NH-CO-Q¹, acylating an amine of formula III



- using an acid of formula HO-CO-Q¹, or an activated derivative thereof;


- (C) for a compound of formula I in which -L¹-Q¹ is -CO-NH-Q¹ and R² is of the form -NH-CO-Q², acylating an amine of formula H₂N-Q¹ using a [1,3]oxazine of formula IV,



- 114 -

wherein Q^2 represents Q^{2B} , Q^{2C} or Q^{2D} ;

- (D) for a compound of formula I in which R^2 is $-L^{2A}-Q^{2A}$ and D is carbonyl, diacylating a compound of formula II using an anhydride of formula V;



- (E) for a compound of formula I in which Q^1 is Q^{1A} and R^a is hydroxy or in which Q^1 is Q^{1B} and R^d is hydroxy, adding hydroxylamine to a corresponding nitrile to afford the compound with an amino(hydroxyimino)methyl group;

- (F) for a compound of formula I in which Q^1 is Q^{1A} and R^a , R^b and R^c are each hydrogen, or in which Q^1 is Q^{1B} and R^d is hydrogen, hydrogenolyzing the N-O bond of a corresponding compound of formula I in which R^a is hydroxy or in which R^d is hydroxy;

- (G) for a compound of formula I in which Q^1 is Q^{1A} and R^a , R^b and R^c are each hydrogen, or in which Q^1 is Q^{1B} and R^d is hydrogen, substituting the methylthio group of a corresponding compound bearing a $-C(SCH_3)=NH$ group with an amino group by treatment with ammonia, or a solvate or salt thereof;

- (H) for a compound of formula I in which Q^1 is Q^{1A} and R^a and R^b together form a methylene or oxo group, cyclizing a corresponding 4-aminomethyl-3-cyanophenyl or 4-aminooxo-3-cyanophenyl compound; or

- (I) for a compound of formula I in which Q^1 is Q^{1A} and R^c is hydroxy, hydrogenolyzing the N-O bond of a corresponding 3-amino-1,2-benzisoxazol-7-yl compound;

- 115 -

whereafter, for any of the above procedures, when a functional group is protected using a protecting group, removing the protecting group; and

5 whereafter, for any of the above procedures, when a pharmaceutically acceptable salt of a compound of formula I is required, it is obtained by reacting the basic form of a basic compound of formula I with an acid affording a physiologically acceptable counterion or the acidic form of an acidic compound of formula I with a base affording a
10 physiologically acceptable counterion or by any other conventional procedure; and

 wherein, unless otherwise specified, L¹, Q¹, R², A³, A⁴, A⁵ and A⁶ have any of the values defined in Claim 10.

15 22. The use of a factor Xa inhibiting compound of formula I substantially as hereinbefore described with reference to any of the Examples.

20 23. A novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.

25 24. A process for preparing a novel compound of formula I substantially as hereinbefore described with reference to any of the Examples.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13424

| A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC | | | | |
|---|--|---|---|--|
| B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/330, 379, 416, 447, 616, 620; 546/234; 548/482; 549/69; 564/155, 157, 164 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE | | | | |
| C. DOCUMENTS CONSIDERED TO BE RELEVANT | | | | |
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. | | |
| Y | US 4,514,416 A (FUJII et al.) 30 April 1985, see entire document, especially column 51, example 42. | 1-4, 10-13, 19, 20 | | |
| Y | US 5,518,735 A (STURZEBECKER et al.) 21 May 1996, column 3. | 1-4, 10-13, 19, 20 | | |
| A | WALLIS R.B., Inhibitors of coagulation factor Xa: from macromolecular beginnings to small molecules. Current Opinion in Therapeutic Patents. August 1993, Vol. 3, No. 8, pages 1173-1179. | 1-4, 10-13, 19, 20 | | |
| <input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex. | | | | |
| <table border="0"> <tr> <td> * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed </td> <td> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family </td> </tr> </table> | | | * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family |
| * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed | "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family | | | |
| Date of the actual completion of the international search 04 SEPTEMBER 1998 | | Date of mailing of the international search report 16 OCT 1998 | | |
| Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230 | | Authorized officer <i>Evelyn Huang</i> EVELYN HUANG Telephone No. (703) 308-1235 | | |

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13424

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5-9, 14-18, 22-24
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13424

A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

A61K 31/38, 31/40, 31/42, 31/165, 31/445; C07C 233/64; C07D 209/44, 211/32, 333/06

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

514/330, 379, 416, 447, 616, 620; 546/234; 548/482; 549/69; 564/155, 157, 164

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-24 in part, drawn to a compound of formula I wherein A3, A4, A5, A6 complete a substituted benzene.

Group II, claim(s) 1-24 in part, drawn to a compound of formula I wherein one of A3, A4, A5, A6 is N.

Group III, claim(s) 1-24 in part, drawn to a compound of formula I wherein two adjacent residues of A3, A4, A5, A6 together form S.

Group IV, claims 1-24 in part, drawn to a compound of formula I wherein two non-adjacent residues of A3, A4, A5, A6 are each N.

Group V, claims 1-24 in part, drawn to a compound of formula I wherein A3 and A4 together form a fused benz ring and A5 and A6 together form -NH-.

The inventions listed as Groups I-V do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: group I is drawn to a substituted benzene, group II is drawn to a pyridine compound, group III is drawn to a thiophene compound, group IV is drawn to a pyrimidine or a pyrazine compound while group V is drawn to an indole compound. A benzene, a pyridine, a thiophene, a pyrimidine or a pyrazine and an indole would not have been of sufficient similarity to allow for a Markush grouping to exhibit utility, absent some teaching of equivalence in the prior art.